

# Northumbria Research Link

Citation: Cornelissen, Katri, Bester, Andre, Cairns, Paul, Tovee, Martin and Cornelissen, Piers The influence of personal BMI on body size estimations and sensitivity to body size change in anorexia spectrum disorders. *Body Image*, 13. pp. 75-85. ISSN 1873-6807

Published by: UNSPECIFIED

URL:

This version was downloaded from Northumbria Research Link: <http://northumbria-test.eprints-hosting.org/id/eprint/46805/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



UniversityLibrary



**Northumbria**  
**University**  
NEWCASTLE

# Northumbria Research Link

Citation: Cornelissen, Katri, Bester, Andre, Cairns, Paul, Tovee, Martin and Cornelissen, Piers (2015) The influence of personal BMI on body size estimations and sensitivity to body size change in anorexia spectrum disorders. *Body Image*, 13. pp. 75-85. ISSN 1873-6807

Published by: Elsevier

URL: <http://dx.doi.org/10.1016/j.bodyim.2015.01.001>  
<<http://dx.doi.org/10.1016/j.bodyim.2015.01.001>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/21521/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

[www.northumbria.ac.uk/nrl](http://www.northumbria.ac.uk/nrl)



# **The influence of personal BMI on body size estimations and sensitivity to body size change in anorexia spectrum disorders**

Katri K. Cornelissen<sup>1</sup>, Andre Bester<sup>2</sup>, Paul Cairns<sup>3</sup>, Martin J. Tovée<sup>4\*</sup> & Piers L. Cornelissen<sup>1</sup>

<sup>1</sup> Department of Psychology, Northumbria University, UK.

<sup>2</sup> Bester Labs, Newcastle upon Tyne, UK.

<sup>3</sup> Department of Computer Science, University of York, UK.

<sup>4</sup> Institute of Neuroscience, Newcastle University, UK.

\*Corresponding Author: Dr Tovée, Institute of Neuroscience, Faculty of Medical Sciences, Newcastle University, NE2 4HH. Email: [m.j.tovee@ncl.ac.uk](mailto:m.j.tovee@ncl.ac.uk)

Running Head: BODY SIZE ESTIMATION IN ANOREXIA SPECTRUM DISORDERS

**Abstract**

In this cross-sectional study, we investigated the influence of personal BMI on body size estimation in 42 women who have symptoms of anorexia (referred to henceforth as anorexia spectrum disorders, ANSD), and 100 healthy controls. Low BMI control participants over-estimate their size and high BMI controls under-estimate, a pattern which is predicted by a perceptual phenomenon called contraction bias. In addition, control participants' sensitivity to size change declines as their BMI increases as predicted by Weber's law. The responses of women with ANSD are very different. Low BMI participants who have ANSD are extremely accurate at estimating body size and are very sensitive to changes in body size in this BMI range. However, as BMI rises in the ANSD participant group, there is a rapid increase in over-estimation concurrent with a rapid decline in sensitivity to size change. We discuss the results in the context of signal detection theory.

**Keywords:** Anorexia Nervosa, Anorexia Spectrum Disorders, Body Size Over-estimation, Signal Detection Theory.

Anorexia nervosa (AN) is a serious psychological and physiological condition, which occurs predominantly in the female population. Current therapeutic regimes have only a limited success in treating this condition (Treasure, Claudino, & Zucker 2010), where the long-term mortality rate has been estimated to be as high as 10% (Berkman, Lohr, & Bulik, 2007). To be able to treat this condition more effectively, we need a better understanding of its central features. Diagnostic criteria for AN include a distorted evaluation of personal body size (DSM-V, 2013), and this is also a key element of psychological models of the disorder (Cash & Deagle, 1997; Fairburn, Cooper, & Shafran, 2003). Body image distortion has been shown to be one of the most persistent of all the eating disorder symptoms, the severity of which seems to predict the long term outcome for patients (Fairburn, Cooper, & Shafran, 2003; Pike, 1998). Furthermore, persistence of body image distortion has been shown to predict the rate of relapse (Channon & DeSilva, 1985; Slade & Russell, 1973) which has been estimated to be as high as 35% (Casper, Halmi, Goldberg, Eckert, & Davis, 1979). While there is evidence to suggest that women with AN under-estimate their body size (Meerman, 1983), or even show performance in size estimation tasks equivalent to non eating-disordered controls (Fernández, Probst, Meerman, & Vandereycken, 1994; Meermann, 1983), most studies have found that patients with AN overestimate their body size (Gardner & Bokenkamp, 1996; Probst, Vandereycken, Van Coppenolle, & Pieters, 1998; Slade & Russell, 1973; Tovée, Benson, Emery, Mason, & Cohen-Tovée, 2003). The disturbance in body size estimation is thought to comprise two components; a perceptual/sensory component and an attitudinal/cognitive component (Cash & Deagle, 1997). The perceptual component is described as an inability to accurately estimate body size. The attitudinal component of body image disturbance consists of dissatisfaction with body shape combined with negative attitudes to weight and shape. Moreover, there is evidence that these effects

1  
2  
3  
4 may be specific to judgements about bodies, and do not generalize to other objects such as  
5  
6 vases (McCabe, Ricciardelli, & Ridge, 2006; Slade & Russell, 1973).  
7  
8

9  
10 Classical psychophysics has been used to try and separate the contributions to body  
11 size estimation made by perceptual ‘sensory’ factors (in this case, the smallest change in  
12 body shape that the participant can detect, indexed by the difference limen, DL) and  
13  
14 attitudinal ‘non-sensory factors’ (the subjective body size criterion, or bias, adopted by the  
15 participant, indexed by the point of subjective equality, PSE). For example, using the method  
16 of constant stimuli, in combination with the video distorting technique, Gardner and  
17  
18 Bokenkamp (1996) reported that women with AN were more likely to over-estimate their  
19 size than non-eating disordered controls, as indexed by a higher PSE on average. On the other  
20 hand, analyses of the same data showed that the smallest difference in stimulus size (DL) that  
21 anorexic participants could reliably detect was no different from controls (i.e., both female  
22 controls and women with AN were equally sensitive at discriminating between different sized  
23 versions of their bodies). Because of this dissociation between PSE and DL, Gardner and  
24  
25 Bokenkamp (1996) suggested that body size over-estimation is entirely due to attitudinal,  
26 non-sensory factors (see also: Gardner & Moncrieff, 1988; Mussap, McCabe, & Ricciardelli,  
27  
28 2008).  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Recently, however, Cornelissen et al. (2013) came to a different conclusion. They re-  
45 analysed a previous study in which women with AN and controls were asked to estimate their  
46 own body size by manipulating an image of themselves using a body morphing program  
47 (Tovée, Benson, Emery, Mason, & Cohen-Tovée, 2003). The software allowed participants  
48 to match their perception of the size of their individual body parts with what they saw on  
49 screen by manipulating slider controls. These had the effect of changing the width and shape  
50 of those body parts. It is possible to calculate the BMI of these self-manipulated bodies from  
51 their perimeter area ratios (Cornelissen, Johns, & Tovée, 2013). As a result, Cornelissen et al.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 could compare directly the participants' estimates of their own BMIs with their actual BMIs.  
5  
6 They found that the inaccuracies in body size estimation could largely be explained by a  
7  
8 known perceptual error in magnitude estimation called contraction bias (Poulton, 1989).  
9  
10 Critically, as shown in Fig. 1A, the relationship between estimated BMI and actual BMI  
11  
12 appeared to be exactly the same for participants with anorexia and controls; there were no  
13  
14 differences in the pattern of contraction bias between the two groups. This figure is a  
15  
16 schematic representation of the results from Cornelissen et al. (2013) in which women with  
17  
18 AN and controls used an interactive software program to estimate body size. The line of  
19  
20 equality (i.e., perfect accuracy) is shown by the dotted black line. The control participants  
21  
22 (whose response distribution is indicated by the cross-hatched region) varied in BMI between  
23  
24 14.7 and 36.8 and the women with AN (indicated by the gray region) varied in BMI between  
25  
26 11.5 and 18.4. The solid black line represents the regression of estimated BMI on actual BMI  
27  
28 and has the same slope and intercept for women with AN and controls.  
29  
30  
31  
32  
33

34           Contraction bias arises when one uses a standard reference or template for a  
35  
36 particular kind of object against which to estimate the size of other examples of that object.  
37  
38 The estimate is most accurate when estimating the size of an object of a similar size to the  
39  
40 reference, but becomes increasingly inaccurate as the magnitude of the difference between  
41  
42 the reference and the object increases. When this happens, the observer estimates that the  
43  
44 object is closer in size to the reference than it actually is. As a result an object smaller in size  
45  
46 than the reference will be over-estimated and an object larger will be under-estimated. Thus,  
47  
48 if we use a “reference body” based on an average of all the bodies we have seen in our life to  
49  
50 make our judgements of body size (Winkler & Rhodes, 2005), individuals with very thin  
51  
52 bodies will over-estimate their own body size, and individuals with very large bodies will  
53  
54 under-estimate their body size (illustrated in Fig 1A). An earlier study by Kuskowska-Wolk  
55  
56 and Rössner (1989) reported results on self-estimation of body size that is also consistent with  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 a contraction bias explanation. In addition, Cornelissen et al. (2013) also found an  
5  
6 independent, modulating effect of psychological factors, as illustrated in Fig. 1A. The thick  
7  
8 dashed black line represents the increase in intercept for the regression of estimated BMI on  
9  
10 actual BMI as psychological concern about body shape and weight increase (i.e., the  
11  
12 regression line moves up the Y-axis as concerns increase). The contraction bias explanation  
13  
14 predicts that for both controls and women with AN, the accuracy of their body size estimation  
15  
16 will be driven by the BMI of the participants. Cornelissen et al. report that this is the case, but  
17  
18 the BMI values of the women with AN in the study by Tovée et al. (2003) all actually fell  
19  
20 within a relatively narrow range 6.9 BMI units (11.5 to 18.4). Most of the variation in BMI in  
21  
22 this study is based on the responses of the control participants who ranged in BMI between  
23  
24 14.7 and 36.8 (22.1 BMI units). Cornelissen et al.'s (2013) assumption is that with a wider  
25  
26 BMI range (including recovering patients to expand the range), the responses of the women  
27  
28 with AN should follow the same pattern as the control participants. This is illustrated in  
29  
30 Figure 1B, where the white arrow shows how the regression of estimated BMI on actual BMI  
31  
32 in these women should track up along the same regression line as in Fig. 1A when BMI  
33  
34 increases. In short, this model predicts that as BMI increases in women with AN, so body  
35  
36 size over-estimation should *decrease*.

37  
38  
39  
40  
41  
42  
43  
44 Alternatively, it is entirely possible that psychological factors represent a stronger  
45  
46 driving force behind body size over-estimation in women with AN than they do for controls.  
47  
48 If so, this could lead to a different outcome. Consistent with this possibility, an individual's  
49  
50 body size (as indexed by BMI) is known to be strongly correlated with body dissatisfaction  
51  
52 (Gardner, Brown, & Boice, 2012; Stice & Shaw, 2002; Striegel-Moore et al., 2004). Women  
53  
54 with AN who have achieved a very low BMI might be expected to have relatively low body  
55  
56 size concerns, but during the recovery process as their weight increases, their body size  
57  
58 concerns would rise in parallel. Therefore, an alternative outcome for women with AN is that  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 as their weight increases, there is a rapid *rise* in the degree of body size over-estimation  
5  
6 reflecting their accelerating concerns about body shape and weight. This is illustrated in Fig.  
7  
8 1C, where the white arrow shows how the regression of estimated BMI on BMI for women  
9  
10 with AN should follow the trajectory of the thick dashed black line.

11  
12  
13  
14 To determine which outcome predicts the accuracy of body size estimations for  
15  
16 women with AN, we have recruited a group of women who have all had a diagnosis of AN,  
17  
18 but who now show significant variation in their BMI and psychological concerns. These  
19  
20 women should more correctly be referred to as suffering from anorexia spectrum disorders  
21  
22 (ANSD). By deliberately taking advantage of individual differences in this way, we can  
23  
24 determine whether the accuracy of body size estimation increases or decreases as the BMI of  
25  
26 ANSD participants varies and therefore determine the relative importance of psychological  
27  
28 factors and perceptual bias as illustrated in Fig. 1B and C.  
29  
30  
31

### 32 33 **Method**

34  
35  
36 The experimental procedures and methods for participant recruitment for this study  
37  
38 were approved by: the local ethics committee at Northumbria University; the Beating Eating  
39  
40 Disorders Organisation (BEAT) and the Northern Initiative on Women and Eating (NIWE)  
41  
42 Organisation.  
43  
44

### 45 46 **Participants**

47  
48  
49 We recruited 42 participants into the study all of whom who have had a formal  
50  
51 diagnosis of anorexia nervosa ( $n=34$ ) according to DSM-IV-R or DSM-5 (American  
52  
53 Psychiatric Association, 2002, 2013) or bulimia nervosa followed by the onset of anorexia  
54  
55 ( $n=8$ ) - it is not uncommon for patients to move between diagnostic categories (Fairburn &  
56  
57 Harrison, 2003). Table 1 shows the details of the participants divided into 3 subgroups:  
58  
59  
60  
61  
62

1  
2  
3  
4 ANSD1: women with a diagnosis of AN who were being treated at the time of testing; ANSD  
5  
6 2: women with a diagnosis of AN who were no longer being treated; AN 3: women with a  
7  
8 diagnosis of either atypical AN ( $n=3$ ) or Bulimia that switched to AN ( $n=8$ ), 4 of whom were  
9  
10 being treated at the time of testing. The right hand columns of Table 1 show the output of  
11  
12 pairwise comparisons of the three subgroup means, adjusted for multiple comparisons, using  
13  
14 the permutation method in PROC MULTEST (SAS v9.3). We acknowledge that this design  
15  
16 is deliberately intended to look at the effect of individual differences within the anorexia  
17  
18 spectrum, as opposed to a design in which we might compare directly between distinct eating  
19  
20 disordered subgroups defined a priori.  
21  
22  
23  
24

25  
26 In addition, 100 participants were recruited from the population of undergraduate  
27  
28 students at Newcastle and Northumbria Universities and from the general population in and  
29  
30 around the Newcastle upon Tyne area, all of whom consented to take part in the study as non-  
31  
32 eating disordered controls. No control participants had a history of eating disorders. The  
33  
34 participants displayed variation in body mass index (BMI), attitudes to eating, body shape  
35  
36 and size, levels of depression and anxiety. For the characteristics of the control participants  
37  
38 see Table 2.  
39  
40  
41

## 42 **Measures**

### 43 **Psychometric measurements**

44  
45  
46 To assess participants' attitudes towards body shape, weight and eating we used: (1)  
47  
48 The Eating Disorders Examination Questionnaire (EDEQ), which is a self-report version of  
49  
50 the Eating Disorder Examination (EDE) structured interview (Fairburn & Beglin, 1994). This  
51  
52 is commonly used as a screening questionnaire for eating disordered behaviour and has been  
53  
54 normed for young women (Mond, Hay, Rodgers, & Owen, 2006). The questionnaire contains  
55  
56 four subscales reflecting the severity of aspects of the psychopathology of eating disorders;  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 the Restraint Scale investigates the restrictive nature of the eating; the Eating Concern scale  
5  
6 to the preoccupation with food and social eating; the Shape Concern subscale investigates  
7  
8 dissatisfaction with body shape and the Weight Concern subscale assesses dissatisfaction  
9  
10 with body weight. The EDE-Q (range 0-6) also measures overall disordered eating behaviour.  
11  
12 Furthermore, it provides frequency data on key behavioural features of eating disorders; (2)  
13  
14 The Eating Attitudes Test (EAT, range 0-78; Garner & Garfinkel, 1979) which investigates  
15  
16 eating habits and dissatisfaction with own body weight and shape. It is a subjective index of  
17  
18 the symptoms displayed by individuals with eating disorders and the test is used as a  
19  
20 screening questionnaire for eating disorders; (3) the 16-item Body Shape Questionnaire  
21  
22 (BSQ, range 0-96; Evans & Dolan, 1993) which indexes the degree of preoccupation and  
23  
24 negative attitude toward body weight and body shape. The participants' level of depression  
25  
26 was measured using the Beck Depression Inventory (BDI, range 0-63; Beck et al., 1961).  
27  
28 Their self-esteem was indexed using the Rosenberg Self-Esteem Scale (RSE, range 0-30;  
29  
30 Rosenberg, 1965).  
31  
32  
33  
34  
35  
36

### 37 **Anthropometric measurements**

38  
39  
40 The participants' body mass index (BMI) was measured with a set of calibrated scales  
41  
42 and a stadiometer. The participants' waist, hip, under-bust and bust circumferences were  
43  
44 measured using measuring tapes. Participants' percentage body fat was measured from  
45  
46 skinfold thickness obtained with Harpenden Callipers. The skinfold measurements were  
47  
48 obtained from eight sites, allowing the percentage of body fat to be calculated (Durnin &  
49  
50 Womersley, 1974). The skinfold measurements were made by an ISAK (International Society  
51  
52 for the Advancement of Kinanthropometry) trained and experienced investigator.  
53  
54  
55  
56

### 57 **Psychophysical measurements**

1  
2  
3  
4 In this study we apply classical psychophysical methods (cf Gardner, 1996) to  
5  
6 measure two components of the participants' judgements of body size: (i) the point of  
7  
8 subjective equality (PSE) and (ii) the difference limen (DL). The PSE is the participant's  
9  
10 estimate of their body size. The DL is an estimate of how sensitive a participant is to changes  
11  
12 in body size and equates to the smallest difference in body size that she can detect. To obtain  
13  
14 these measurements, we use the method of constant stimuli in a two alternative forced choice  
15  
16 paradigm. On every trial of the task, the participants are presented with an image of a woman  
17  
18 and they have to respond by button press whether they believe the image to be smaller or  
19  
20 larger than themselves. The BMI of the body varies on each presentation from smaller than to  
21  
22 larger than the actual BMI of the participant. For each participant, we then plotted the  
23  
24 proportion of 'larger' responses (y-axis) as a function of the BMI of the stimulus (x-axis), and  
25  
26 fitted a cumulative Gaussian (sigmoid) curve to the data, as illustrated in Fig. 3 D. The PSE is  
27  
28 then defined from this curve as the BMI at which participants would respond 'larger' 50% of  
29  
30 the time. Therefore, the PSE corresponds to the midpoint of the curve where, effectively,  
31  
32 participants would be equally like to respond 'smaller' or 'larger' because a stimulus with  
33  
34 this BMI has the same size as they believe themselves to have. The value of BMI at the PSE  
35  
36 may be smaller (i.e., an underestimation), the same as or larger (an over-estimation) than a  
37  
38 participant's actual BMI. The DL is the difference in the BMI of the stimuli falling between  
39  
40 the 25% and 75% 'larger' response points (see Gescheider, 1997). This range captures the  
41  
42 steepness of the psychometric curve. Participants who are very sensitive to small changes in  
43  
44 body size will have a steeper psychometric function with a correspondingly small DL. It is  
45  
46 important to emphasize that there is no necessary correlation between PSE and DL.  
47  
48 Participants who over-estimate or under-estimate their body size may be equally likely to be  
49  
50 sensitive (small DL) or insensitive (large DL) to changes in body size.  
51  
52  
53  
54  
55  
56  
57  
58  
59

### 60 **Stimulus image preparation**

1  
2  
3  
4           Creating stimulus images which correctly represent how body shape changes as a  
5  
6 function of changing BMI is difficult, because these changes are highly non-linear. One  
7  
8 method that has been used previously is the video-distortion technique (VDT) (see e.g.,  
9  
10 Gardner and Bokenkamp, 1996; Probst, Vandereycken, & Van Coppenolle, 1997) in which  
11  
12 2D images of people are stretched or compressed in the horizontal dimension. However, this  
13  
14 linear method is problematic as it creates shape changes particularly in the shoulder and hip  
15  
16 regions which tend to be unrealistic (see below). Alternative methodologies can be used, such  
17  
18 as morphing between images of high and low BMI bodies. While this is an improvement on  
19  
20 the VDT in principle, because it is a non-linear method, it is nevertheless extremely difficult  
21  
22 to maintain the combination of high feature definition and stable identity of the person in the  
23  
24 morphed images across a wide range of BMI values. Inevitably, some form of averaging or  
25  
26 smoothing is required which reduces the realism in the resultant images. For these reasons,  
27  
28 we instead used film industry computer-generated imagery (CGI) methods to create graded  
29  
30 3D images of individuals.  
31  
32  
33  
34  
35  
36

37           Figure 2 illustrates the comparison between the VDT and CGI methods. The top row  
38  
39 shows three CGI images from our database of stimuli. From left to right, they represent  
40  
41 approximately BMIs 13, 17 and 21 respectively. The second row shows the body shape  
42  
43 changes produced by applying the video distortion technique (VDT) to the central image in  
44  
45 the first row. The horizontal compression (left image) and expansion (right image) for the  
46  
47 images on the second row were made so that waist widths corresponded to the BMI 13 and  
48  
49 21 images from the top row, respectively. The third row shows the difference between the  
50  
51 two methods of image manipulation. The grey shaded outline represents the CGI image and  
52  
53 the dotted line the VDT image for BMI 13 (left) and BMI 21 (right). Row three shows how  
54  
55 for thinner images, the VDT compresses the width of the hips and shoulders more than the  
56  
57 CGI technique. In comparison, for larger images, the VDT exaggerates the width of the hips  
58  
59  
60  
61  
62  
63  
64  
65

and shoulders relative to the CGI method. In addition, the width of the gap between the thighs changes in opposite directions: consistent with reality, the CGI method shows a reduction in this gap with increasing BMI, whereas the VDT shows an increase.

Particular advantages of the CGI method are: i) the identity of the person in the image is clearly maintained over a wide BMI range; ii) the body shape changes at different BMI levels are extremely realistic and iii) the 3D rendered stimulus images are high definition and photorealistic. Another critical feature is that the stimuli are calibrated for BMI. We used the Health Survey for England (HSE 2003 & 2008) datasets to create calibration curves between waist and hip circumferences and height derived from ~5000 females in the UK, aged between 18 and 45. Because our CGI models exist in an appropriately scaled 3D world, having set the height of our models (1.6m) we can therefore measure their waist and hip circumferences, and compare these with our HSE calibration curves in order to compute their BMI (Cornelissen, Toveé, & Bateson, 2009). Finally, we carried out two further qualitative checks of the plausibility of the overall body shape changes that we created in the modelling environment. First, we compared the 3D volumes and shapes from our CGI modelled bodies to a 3D statistical model of the relationship between BMI and shape changes in 114 real bodies (Hasler, Stoll, Sunkel, Rosenhahn, & Seidel, 2009). Secondly, we compared our images against a library of digital photographs of 220 women in a standard pose who varied in BMI from 11 (emaciated) to 45 (obese) (Tovéé, Benson, Emery, Mason, & Cohen-Tovéé, 2003).

## Procedure

For the presentation of stimuli, we used a Python program written by one of the authors to implement the two alternative forced choice task. Stimuli were presented on an 18" flat panel LCD screen (1280w x 1024h pixel native resolution, 32-bit colour depth) for as

long as it took participants to make a decision. At the standard viewing distance of ~60cm, the CGI model subtended ~26° vertically and ~8° degrees horizontally. Each participant first judged 7 images covering the whole BMI range (from 12.5 to 44.5 in equal BMI steps) presented in 2 separate blocks. Each stimulus image appeared 10 times in each block, and the order of presentation was randomized. Based on the responses from each block, the participants' point of subjective equality or PSE (the BMI they believe themselves to be) was calculated automatically by fitting a cumulative normal distribution. These two values were then averaged to give an initial estimate of the participant's PSE. On the basis of this initial estimate, the program presented a further set of 21 images (spread over a range of 5 BMI units centered on the participant's initial PSE, at a spacing of 0.25 units per image) for the participants to judge. Each image was presented 10 times in randomized order. This final set of judgements allowed us to calculate a definitive estimate of PSE as well as the difference limen or DL (that is how sensitive participants are to changes in BMI).

## Results

### Task reliability

The responses to the questionnaires across the sample showed good internal reliability scores. For RSE, BDI, BSQ, EAT and EDEQ, Cronbach's alpha was: .89, .93, .96, .92 and .96 respectively. For the PSE scores from the 2AFC task, we estimated reliability in two ways. First, for the entire sample, we divided the last block of body size estimation trials into a first and a second half (i.e., in time), in order to estimate split-half reliability. We carried out separate probit analyses on each half of the data to obtain a PSE estimate from each half. The correlation between the two sets of PSE scores ( $r=.92, p<.0001$ ) suggests good reliability. In addition, we asked 17 participants to return to the laboratory to re-run the body size judgement task. The shortest interval for these test retest measurements was ~ 1 month.



The Pearson correlation between PSE estimates was  $r=.93$  ( $p<.0001$ ) and a dependent samples t-test comparison of the means was not significantly different from zero ( $M = 0.48$ ,  $p=.333$ ).

### **ANSD participants**

As Table 1 shows, there are significant differences between the three subgroups of ANSD participants on some psychometric measures. For example, consistent with previous studies (e.g., Tovée et al. 2000, 2003), the group who are currently under treatment (i.e., ANSD1) have significantly lower BMIs, lower self-esteem, higher tendency towards depression and the most disturbed attitudes to eating compared to the other two groups (ANSD2 and ANSD3). This raises the question whether the relationships between PSE, DL and BMI in the three ANSD subgroups might be different. If so, then the distinction between the three groups should be maintained when comparing them statistically to healthy controls. Alternatively, if, for example, there are no statistically significant differences in the relationships between PSE and BMI for these three subgroups, we would gain statistical power in our analyses by treating all participants with a history of AN as one group. Therefore, we used PROC MIXED (SAS v9.3) to test whether there is a main effect of ANSD grouping on the regressions of PSE on BMI, and separately, DL on BMI. Since the distribution of raw DL scores was non-normal (Shapiro-Wilk's  $W=.62$ ,  $p < .001$ ) we applied a logarithmic transformation to these data. For PSE, the mixed linear model explained 71% of its variance. The Type III test of the fixed effect of BMI was significant at the specified .05 level,  $F(1,37) = 67.36$ ,  $p < .001$ , but the tests for ANSD group,  $F(2,37) = 0.55$ ,  $p = .580$ , and the interaction between BMI and ANSD group,  $F(2,37) = 0.54$ ,  $p = .587$ , were not. For DL, the mixed linear model explained 15.8% of its variance. The Type III test of the fixed effect of BMI was significant at the specified .05 level,  $F(1,37) = 7.11$ ,  $p = .011$ , but the tests for

1  
2  
3  
4 ANSD group,  $F(2,37) = 1.09, p = .347$ , and the interaction between BMI and ANSD group,  
5  
6  $F(2,37) = 1.07, p = .354$ , were not. Since we could find no statistically robust effects of the  
7  
8 subgroups in these preliminary analyses, henceforth we treat all participants with ANSD as  
9  
10 one group.  
11

### 12 13 14 **Univariate statistics**

15  
16  
17 Table 2 shows the means and standard deviations (SD) for the participant  
18  
19 characteristics, separated according to whether they belong to the anorexia spectrum  
20  
21 disorders (ANSD) or the non eating-disordered control group. The right most column of  
22  
23 Table 2 shows the output of pairwise comparisons of these group means, adjusted for  
24  
25 multiple comparisons, using the permutation method in PROC MULTTEST (SAS v9.3). For  
26  
27 The Eating Disorder Examination Questionnaire, a global eating disorder score was compiled  
28  
29 across the subtests which combines the evaluation of restrictive eating behaviour, participant  
30  
31 psychological concerns about eating, concerns about their own weight and concerns about  
32  
33 their own body shape. Consistent with previous literature we found that when compared to  
34  
35 non-eating disordered controls, the ANSD participants had statistically significantly lower  
36  
37 BMIs and percentage body fat, elevated concerns about body shape, eating behaviour and  
38  
39 body weight (EAT, BSQ, EDEQ), greater tendency towards depression, reduced self-esteem  
40  
41 and significantly greater over-estimation of body size (OE). The relatively high mean BMI  
42  
43 for the ANSD participants reflects that fact that an adequate test of our hypotheses *requires*  
44  
45 this participant group to have a wide BMI range. Finally, we found no statistically significant  
46  
47 differences between the group means for the sensitivity measure for body size estimation  
48  
49 (DL) or chronological age.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

### 60 **Multivariate statistics**

We used PROC MIXED (SAS v9.3) to build a mixed linear model with estimated BMI (i.e., PSE) as the outcome variable, and actual BMI and GROUP (i.e., ANSD versus healthy controls) as predictor variables. In addition, we wanted to control for any influence of AGE and the psychometric variables (RSE, BDI, BSQ, EAT and EDEQ). In order to avoid the possibility of introducing substantial variance inflation, we first checked for evidence of co-linearity amongst the psychometric variables. Across the sample of 142 participants the Pearson correlations between RSE and BDI, RSE and BSQ, RSE and EAT, RSE and EDEQ, BDI and BSQ, BDI and EAT, BDI and EDEQ, BSQ and EAT, BSQ and EDEQ, and EAT and EDEQ were: -.75, -.61, -.62, -.57, .60, .66, .56, .72, .84 and .79 respectively. All correlations were statistically significant at  $p < .001$ . Given these substantial correlations, we therefore used PROC FACTOR in SAS v9.3 (SAS Institute, North Carolina, US) to carry out an iterated principal factor analysis with rotation in order to identify the significant latent variable(s) in the psychometric data. We then used the factor scores from these latent variable(s) in our statistical models. The Kaiser-Meyer-Olkin (*KMO*) measure of sampling adequacy (which indicates the degree of diffusion in the pattern of correlations) was .81 suggesting an acceptable sample. One factor had an Eigen value greater than Kaiser's criterion of 1 (i.e., 3.39) which explained 98% of the variance. The scree plot showed an inflexion, i.e., Cattell's criterion which also justified retaining just the one factor. The residuals were all small, and the overall root mean square off-diagonal residual was 0.081, indicating that the factor structure explained most of the correlations. The factor loadings for RSE, BDI, BSQ, EAT and EDEQ were: -.76, .76, .86, .86, .86 respectively. This latent variable, referred to henceforth as PSYCH, represents a combination of the attitudes thought to contribute to body size disturbance: disturbed attitudes to eating, weight, and shape, and low self-esteem and depression.

**PSE**

The mixed linear model for PSE explained 71% of its variance. Both ANSD and control participants showed a positive, linear relationship between actual BMI and PSE,  $F(1,137) = 207.72, p < .001, 95\% CI [0.68, 0.90]$ . Critically, however, the pattern of estimation by women with ANSD is statistically different from that of the controls because we found a significant main effect of GROUP,  $F(1,137) = 14.90, p < .001, 95\% CI [-19.08, -6.16]$  as well as an interaction between GROUP and BMI,  $F(1,137) = 15.33, p < .001, 95\% CI [.29, .89]$ .

Figure 3A illustrates the outcome from the mixed linear modelling for PSE. White dots with solid white regression line and gray triangles with solid black regression line represent participants with ANSD and non eating-disordered controls respectively. The dotted black diagonal line in Fig. 3 A shows the line of equality (i.e., perfect estimation) between actual BMI (x-axis) and estimated BMI (y-axis) as indexed by PSE. Data points above this line represent over-estimation, while data points below it represent under-estimation. Fig. 3A therefore shows that low BMI control participants tend to over-estimate body size; control participants whose BMI was in the mid-range were reasonably accurate, and high BMI control participants tended to under-estimate body size. The regression slope for this relationship was significantly *less* than 1,  $\beta = 0.82, F(1,97) = 9.96, p < .005$ , and is therefore consistent with contraction bias in the controls. Participants with ANSD, however, showed a very different pattern of results. Low BMI participants with ANSD were the most accurate at body size estimation. Thereafter, in participants with higher BMI, individuals with ANSD systematically over-estimated body size, and the magnitude of this over-estimation increased in direct proportion to increasing BMI. The regression slope for PSE on BMI in ANSD participants was significantly *greater* than 1,  $\beta = 1.39, F(1,39) = 5.86, p < .05$ . Clearly, this is not consistent with contraction bias, and therefore these results do not support

1  
2  
3  
4 hypothesis one as outlined in Fig. 1 A, but do support hypothesis 2. Finally, we found a  
5  
6 significant Type III test of the fixed effect of PSYCH,  $F(1,137) = 17.23, p < .001, 95\% CI$   
7  
8  $[0.59, 1.66]$  on PSE. However, we found no evidence for any two- or three-way interactions  
9  
10 involving PSYCH. This result is illustrated in Fig. 3 B. White and black regression lines  
11  
12 represent participants with ANSD and non eating-disordered controls respectively. Dashed  
13  
14 and solid lines represent ‘high’ (+1 SD) and ‘low’ (- 1 SD) effects of PSYCH respectively.  
15  
16 This result suggests that any participant with a greater tendency towards depressive  
17  
18 symptoms, shape and weight concern, *irrespective of group*, was more likely to over-estimate  
19  
20 their body size. Therefore, while the significant interaction between GROUP and BMI  
21  
22 supports hypothesis 2, it is unlikely that this can be attributed to psychological factors alone,  
23  
24 because any influence of PSYCH on PSE was essentially the same for ANSD and healthy  
25  
26 controls.  
27  
28  
29  
30

31  
32  
33 How robust is this result? To address this question, we knocked out participants one  
34  
35 by one from the ANSD sample by systematically restricting the upper limit of the sample for  
36  
37 BMI and refitting the model above. We retained a statistically significant effects of GROUP  
38  
39 even when the sample size for the ANSD participants was reduced to 67% ( $n=28$ ) of the  
40  
41 original 42, at a BMI cut-off of 22.3. Therefore, we suggest that our result is indeed robust.  
42  
43 Moreover, this test of the model is considerably more stringent than applying standard  
44  
45 regression diagnostics, because no data point in the analysis of the full sample exceeded the  
46  
47 Cook’s  $D$  or  $DFITS$  critical values for undue influence in the model (i.e., 1 & 2  
48  
49 respectively). As a further check, we also re-ran the whole analysis having removed the 8  
50  
51 participants whose initial diagnosis was bulimia, and obtained remarkably similar,  
52  
53 statistically robust results. For this reduced model, the Type III tests of the fixed effects of  
54  
55 BMI,  $F(1,129) = 175.33, p < .001, 95\% CI [0.68, 0.90]$ , PSYCH,  $F(1,129) = 14.40, p < .001,$   
56  
57  $95\% CI [0.50, 1.60]$ , GROUP,  $F(1,129) = 10.85, p = .001, 95\% CI [-18.03, -4.50]$ , and the  
58  
59  
60  
61  
62

interaction between BMI and GROUP,  $F(1,129) = 11.03$ ,  $p = .001$ , 95% *CI* [0.22, 0.85], were all significant at the specified .05 level.

## DL

The mixed linear model for  $\log_{10}$ DL explained 22.1% of its variance. Both ANSD and control participants showed a positive, linear relationship between actual BMI and  $\log_{10}$ DL,  $F(1,137) = 21.98$ ,  $p < .001$ , 95% *CI* [0.0068, 0.025]. However, the significant main effect of GROUP,  $F(1,137) = 7.05$ ,  $p = .008$ , 95% *CI* [-1.32, -0.19] as well as the interaction between GROUP and BMI,  $F(1,137) = 5.00$ ,  $p = .027$ , 95% *CI* [0.0034, 0.056] show that the pattern of estimation by women with ANSD is statistically different from that of the controls, as can be seen in Fig. 3C. White dots with a solid white regression line and gray triangles with a solid black regression line represent participants with ANSD and non eating-disordered controls respectively. The control participants only showed a modest reduction in sensitivity to the task (i.e., increasing DL) with increasing BMI. This pattern was much more dramatic for participants with ANSD: low BMI participants with ANSD showed much smaller DL values than controls with similar BMI, suggesting very high sensitivity to the task. As the BMI of participants with ANSD increased, DL increased steeply towards the values of control participants, at a BMI of ~26. Finally, we did not find an effect of PSYCH,  $F(1,137) = 0.75$ ,  $p = .389$ , 95% *CI* [-0.026, 0.067] on  $\log_{10}$ DL.

In summary, the results for PSE and DL suggest significant differences in the perceptual aspects of body size estimation when participants with ANSD and non-eating disordered controls are compared. Moreover, the particular pattern of differences in participants with ANSD depends on the BMI of the observer: very accurate and sensitive body size estimation at low BMI, but large over-estimation with low sensitivity at high BMI.

## Discussion

1  
2  
3  
4 The aim of the current study was to investigate the relationship between BMI, body  
5 size estimation and psychological status in women who have had a diagnosis of anorexia  
6 nervosa and compare their responses to those of non eating-disordered controls. Based on  
7 prior literature, we predicted two possible outcomes. According to the contraction bias model  
8 (Fig. 1B), increasing BMI in women with ANSD should lead to *reducing* body size over-  
9 estimation, because their judgements should follow the same trajectory with increasing BMI  
10 as has been shown for non eating-disordered controls. In contrast, based on studies which  
11 report strong positive correlations between BMI/body size and psychological concerns about  
12 body shape and weight, Fig. 1C illustrates an opposing prediction: that body size over-  
13 estimation should *increase* systematically with increasing BMI.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 In our control participants, the accuracy of body size estimation is linearly predicted  
29 by their own BMI, but with a slope less than unity (see Fig. 3A): low BMI controls over-  
30 estimate body size, mid-range BMI controls' estimates are the most accurate and high BMI  
31 controls under-estimate. This pattern of responses is entirely consistent with a normal feature  
32 of magnitude estimation, i.e., contraction bias (Poulton, 1989). Contraction bias assumes that  
33 a body size judgment is made by comparing a body with an internal standard or reference  
34 based on an average of all the bodies that a person has ever seen. This judgement is most  
35 accurate when comparing a body similar in size to the internal reference body, and  
36 increasingly less accurate as the two diverge. In these latter cases, when there is an increasing  
37 difference between the reference and the body to be estimated, the observer tends to select a  
38 response closer to the reference value than it should be, hence the term contraction bias (i.e.,  
39 sizes much larger than the reference are underestimated and smaller sizes than the reference  
40 are overestimated).  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 This contraction bias explanation is consistent with results from other studies. For  
5  
6 example, Tovée et al. (2003) used a large sample of both eating disordered and control  
7  
8 participants and found that as the participants' own BMI declines, their overestimation  
9  
10 increases, with highest overestimation in the very low BMI individuals. Additionally  
11  
12 contraction bias can also explain the fact that previous studies have consistently shown that  
13  
14 obese people under-estimate their size relative to normal BMI people (Kuchler & Variyama,  
15  
16 2003; Kuskowska-Wolk et al., 1989; Maximova et al., 2008; Robinson & Kirkham, 2013;  
17  
18 Truesdale & Stevens, 2008; Wetmore & Mockdad, 2012). Critically, given that contraction  
19  
20 bias is a normal attribute of our perceptual systems, these studies also suggest that there is a  
21  
22 substantial perceptual component to body size estimation that will be dependent on an  
23  
24 individual's own BMI and not on their psychological state.  
25  
26  
27  
28  
29

30 Although contraction bias describes the gradient of the response between the accuracy  
31  
32 of the estimation against the BMI of the body being judged by the controls, the intercept for  
33  
34 this relationship is also influenced by body shape concerns (i.e., the function moves up or  
35  
36 down the y-axis depending on the level of their psychological concerns, illustrated in Fig.  
37  
38 3B). This shows that for a body of a given BMI, the degree of size over- or under-estimation  
39  
40 will also be modulated by the psychological state of the observer, even in the control  
41  
42 participants.  
43  
44  
45  
46

47 The difference limen (DL) for the control participants rises as the participants' own  
48  
49 BMIs rise (i.e., the participants' sensitivity to changing body size declines as their own BMI  
50  
51 rises). This can be explained by another well-established perceptual effect, Weber's law. This  
52  
53 states that the just noticeable difference (JND) between two stimuli will be a constant  
54  
55 proportion of their magnitude, leading to a constant Weber fraction over the stimulus range  
56  
57 (i.e.,  $\Delta I / I = K$ , where  $I$  = stimulus magnitude and  $K$  = constant). This means that, for bodies,  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 it is easier to notice, for example, a one BMI unit difference between two low BMI bodies  
5  
6 than between two high BMI bodies. Over the full range of BMI, discriminating between  
7  
8 higher BMI bodies requires progressively larger differences in BMI between stimuli, which  
9  
10 in turn leads to progressively increasing DL values. Indeed, the data for controls in Fig. 3C  
11  
12 show an excellent fit to Weber's Law because the Weber fractions at, e.g., BMI 20 and 40 are  
13  
14 0.036 and 0.038 respectively, which fall within 5% of each other.  
15  
16

17  
18  
19 In this study, women with ANSD showed a qualitatively different pattern of results  
20  
21 compared to the controls, and these differences are statistically robust. For women with  
22  
23 ANSD, the lowest BMI individuals are actually the most accurate in their body size  
24  
25 estimation and they are also very sensitive to changes in body size (i.e., they show a low DL  
26  
27 value) in low BMI bodies. This sensitivity is considerably greater than that seen for controls.  
28  
29 However, as the ANSD individuals' BMI increases, they start to systematically over-estimate  
30  
31 their body size (as indexed by PSE) and the magnitude of this over-estimation scales linearly  
32  
33 with their BMI. This is reflected in the sharp increase in the DL values, indicating a rapid  
34  
35 reduction in sensitivity to changes in body size as the BMI of the bodies' increases.  
36  
37 Moreover, the gradient of this relationship for DL in women with ANSD far exceeds the  
38  
39 prediction from Weber's law. Overall, therefore, the pattern of over- and under-estimation of  
40  
41 body size is quite different to that described in the control participants, is the opposite of the  
42  
43 pattern of responses predicted by contraction bias, and is more consistent with the prediction  
44  
45 in Fig. 1C.  
46  
47  
48  
49  
50

51  
52 In Fig. 1C, we had originally hypothesised that the low BMI women with anorexia  
53  
54 would have relatively low body size concerns and that the recovering women with anorexia  
55  
56 would show an increase in concerns as their BMI rose. What we found was that women with  
57  
58 ANSD did have greater psychological concerns, on average, than controls (see Table 2).  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 However, the influence of psychological concerns on the relationship between personal BMI  
5  
6 and body size estimation was the same for both groups (see the difference between  
7  
8 continuous and dotted regression lines in Fig. 3B) – i.e., increasing psychological concerns  
9  
10 were linked to higher body size estimations across the entire sample, changing the intercepts,  
11  
12 but not the slopes, by the same amount for both groups. This means that changing  
13  
14 psychological concerns cannot be directly responsible for the perceptual differences in the  
15  
16 two groups that we observe, i.e., the different slopes for PSE and DL in Fig. 3A and C.  
17  
18 Instead, we suggest that the key to explaining our findings lies in the different ways that  
19  
20 women with ANSD *interpret* changes in body size and psychological concern, compared to  
21  
22 controls. In what follows, we use signal detection theory, to account for the results in women  
23  
24 with ANSD, based on the reduction in their sensitivity to changes in body size as the BMI of  
25  
26 the bodies being judged increases.  
27  
28  
29  
30  
31

32  
33 Consider the paradigm being used. Each participant has to say whether a body being  
34  
35 shown is fatter or thinner than themselves. As the BMI of ANSD participants increases, they  
36  
37 become less able to distinguish small differences between the BMI of the stimuli and the  
38  
39 body size they believe they have because they are less sensitive (i.e., increasing DL) to  
40  
41 differences in body size (see Fig. 3C). This means that in order for a low BMI and a higher  
42  
43 BMI woman with anorexia to have the same confidence in expressing a judgement about  
44  
45 their respective body sizes, the participant who has a higher personal BMI, will only response  
46  
47 ‘larger than me’ to stimuli that are, proportionately, considerably larger than they are, as  
48  
49 compared to the participant who has a lower personal BMI. Psychophysically, this is  
50  
51 reflected in a so-called criterion shift: i.e., not only does the slope of the psychometric  
52  
53 function become shallower for the participant with a higher personal BMI (i.e., reduced  
54  
55 sensitivity), but the curve itself is also shifted to the right (see Fig. 3D). Thus, we propose  
56  
57 that because of their reduced sensitivity to changes in body size, the higher BMI ANSD  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 participants shift their response criterion in order to remain confident that their responses are  
5  
6 correct, which in turn causes an increase in PSE and consequent over-estimation of body size.  
7  
8 We suggest that an insistence to maintain the same confidence in the decisions made by  
9  
10 women with ANSD is consistent with their documented aversion to making errors in  
11  
12 judgements (Kaye, Fudge, & Paulus, 2009; Wagner et al., 2007). We ran a post-hoc  
13  
14 correlation analysis which supports this interpretation, because the level of psychological  
15  
16 concern in women with ANSD was positively correlated with body size estimates (i.e., PSE;  
17  
18  $r = .30, p = .05$ ) but independent of sensitivity to body size change (i.e., DL;  $r = .17, p = .29$ ).  
19  
20 When we partialled out the influence of sensitivity (DL), the correlation between  
21  
22 psychological concerns and body size estimation was rendered non-significant ( $r = .25, p =$   
23  
24  $.11$ ). This therefore suggests that sensitivity to change in body size plays a critical role in  
25  
26 modulating the relationship between psychological concern and body size estimation in  
27  
28 women with anorexia.  
29  
30  
31  
32  
33

34  
35 Gardner and Moncrieff (1988) used Signal Detection Theory to analyse their data  
36  
37 from a 2AFC task in which participants had to judge whether images of themselves were  
38  
39 normal or distorted. While they found no difference in task sensitivity (indexed by d-prime)  
40  
41 between women with AN and controls, nevertheless, the women with AN showed a more lax  
42  
43 response criterion. Gardner and Moncrieff (1988) suggested that this may make women with  
44  
45 AN more likely than controls to report an image of themselves as distorted. If we assume an  
46  
47 average height for their sample of 1.6m, then the average BMI of the women with AN and  
48  
49 controls in the Gardner and Moncrieff study were ~17.5 and ~22 respectively. At these BMI  
50  
51 levels, we do see marked differences in task sensitivity (indexed by DL) between women  
52  
53 with ANSD and controls in the current study. We suggest two reasons for the differences  
54  
55 between studies. The first is that Gardner and Moncrieff (1988) used the video distortion  
56  
57 technique (VDT) to produce their body stimuli. As illustrated in Fig. 2, there are systematic  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 differences in the body shapes produced by the VDT compared to our CGI method which  
5  
6 might induce different patterns of responses in women with ANSD and controls. Secondly, it  
7  
8 is possible that the use of a distortion question (i.e., normal versus distorted) may have  
9  
10 invoked a different response pattern from the women with AN than the thinner or fatter  
11  
12 question we used and which is potentially a more psychologically and emotionally charged  
13  
14 judgement for them to make. Our results suggest that women with ANSD are extremely  
15  
16 sensitive to shape change in low BMI bodies and so they may be better placed than controls  
17  
18 to detect visual changes in the VDT images which do not usually accompany weight  
19  
20 loss/gain. Further research is needed to clarify the nature and implications of these  
21  
22 methodological differences.  
23  
24  
25  
26  
27

28         While the account we have given of our data uses ideas from signal detection theory,  
29  
30 it does leave open the question of what mechanism might cause the dramatic differences in  
31  
32 task sensitivity demonstrated by women with ANSD; clearly a specific focus for future  
33  
34 research. One possibility is that the higher sensitivity (i.e., lower DL) to changes in body size  
35  
36 in ANSD women who have low BMI, relative to controls, is due to an expertise effect (i.e.,  
37  
38 the development of expertise in low BMI body discrimination). Women with AN spend a  
39  
40 great deal of time looking at low BMI bodies including their own, but also online as part of  
41  
42 their obsession with the thin ideal (Norris, Boydell, Pinhas, & Katzman, 2006; Ransom, La  
43  
44 Guardia, Woody, & Boyd, 2010). This could explain why our participants were more  
45  
46 sensitive to low BMI body change than the controls. Repeated evaluation and discriminations  
47  
48 of low BMI bodies could allow the development of an expertise in discriminating between  
49  
50 low BMI bodies. ANSD individuals' sensitivity to size change in higher BMI bodies showed  
51  
52 a rapid decline, far faster than is seen in the control participants. Body shape changes in a  
53  
54 non-linear fashion with increasing BMI (Wells, Treleaven, & Cole, 2007), so the pattern of  
55  
56 shape change with weight increase is different in low BMI bodies as compared to heavier  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 BMI bodies. Arguably, therefore, the expertise developed by women with ANSD might be  
5  
6 specific to low BMI bodies and may not generalise to discriminating between higher BMI  
7  
8 bodies.  
9

10  
11  
12         Although we have emphasised the role of perceptual factors in our explanation, we  
13  
14 acknowledge that a range of psychological, cultural and socio-economic factors influence our  
15  
16 judgement of bodies. A number of studies have suggested that we make judgements about  
17  
18 complex stimuli such as faces and bodies by reference to a template based on the average of  
19  
20 all that class of stimuli that they have seen (e.g., Leopold, O'Toole, Vetter, & Blanz, 2001;  
21  
22 Winkler & Rhodes, 2005). This average can be selectively manipulated by showing a  
23  
24 sequence of thin or fat bodies to shift an observer's perception of what constitutes a normal  
25  
26 body (Winkler & Rhodes, 2005). However, this "visual diet" is not the only factor in  
27  
28 preferred body size. For example, showing larger bodies in a positive light can increase the  
29  
30 preference for larger bodies (Boothroyd, Tovée, & Pollet, 2012). An example of this "visual  
31  
32 valency" is the change in body preferences as people move between cultures. People living in  
33  
34 rural KwaZulu Natal in South Africa prefer a much larger BMI (Tovée, Swami, Furnham, &  
35  
36 Mangalparsad, 2006), but people from KwaZulu Natal moving to the UK shift their  
37  
38 preferences towards a lower BMI. The average BMI of people in both regions is not  
39  
40 significantly different, but in KwaZulu Natal a heavier body is associated with health and  
41  
42 higher socioeconomic status whereas in the UK the opposite is the case. Thus, it is the value  
43  
44 placed a particular body size by a culture that seems to predict preferred body size. So  
45  
46 multiple factors influence body preferences.  
47  
48  
49  
50  
51  
52  
53

54         Our results have important implications for treatment of people at both ends of the  
55  
56 weight spectrum. Women with ANSD are very accurate in their body size judgements when  
57  
58 they have a low BMI (BMI<17.5) and tend to actually slightly under-estimate their body size.  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 This suggests that body size over-estimation as a maintenance factor for low BMI women  
5  
6 with anorexia has a comparatively weak effect. However, as they are very sensitive to  
7  
8 changes in the size and shape of a low BMI body, they are also able to detect any increase in  
9  
10 their weight. This may stimulate a renewal of weight reduction behaviours. In this case it is  
11  
12 not an inaccurate perception of their body that is the potential problem, but instead their  
13  
14 ability to accurately detect weight increase.  
15  
16  
17

18  
19 However, as their BMI rises, women with ANSD rapidly develop an over-estimation  
20  
21 of their body size and a rapid decrease in the sensitivity with which they can detect changes  
22  
23 in body size. These changes occur significantly faster as participant BMI increases in women  
24  
25 with ANSD as compared to controls of a corresponding BMI (see Figure 3A & B). This has  
26  
27 important implications for potential relapse in recovering patients. They will see themselves  
28  
29 as larger than they actually are and a large proportion of this over-estimation will be based on  
30  
31 perceptual factors not easily treated through talking therapies such as cognitive behavioural  
32  
33 therapy (CBT). Given the core nature of body size concerns for AN (Cash & Deagle, 1997;  
34  
35 Fairburn, Cooper, & Shafran, 2003), this represents a potentially important factor in any  
36  
37 subsequent relapse and may contribute to the relatively high relapse rate (Channon &  
38  
39 DeSilva, 1985; Slade & Russell, 1973). However, this is not to say that CBT cannot play a  
40  
41 role in reducing body size over-estimation. For a given BMI, our results suggest that the  
42  
43 degree of over-estimation by women with ANSD is modulated by their body size and shape  
44  
45 concerns. So talking therapies such as CBT, can potentially reduce the over-estimation of  
46  
47 body size by targeting these concerns. However, to improve perceptual judgements a  
48  
49 perceptual training program may also be needed to try and recalibrate their judgements.  
50  
51  
52

53  
54  
55  
56 The results also have significant implications for the treatment of obesity. A number  
57  
58 of studies have suggested that levels of obesity are increasing in the general population  
59  
60  
61

(Ogden, Carroll, Curtin, McDowell, Tabak, & Flegal, 2006; Swinburn, Sacks, Hall, McPherson, Finegood, Moodie, & Gortmaker, 2011). At the same time, it has been reported that people who are overweight have difficulty in detecting that their weight is increasing (Truedale & Stevens, 2008; Wetmore & Modkdad, 2012). This has been explained by an adaptation effect, suggesting that the general trend in the population to be heavier has redefined what we judge to be overweight (Robinson & Kirkham, 2013). However, our results suggest two other factors could also play a role. Firstly, contraction bias produces a systematic under-estimation of our body size as we gain weight. Secondly, Weber's law predicts that we will be systematically less able to detect weight increase (as indexed by the JND) as our BMI rises. Both of these factors could act to make it more difficult for overweight people to detect weight increase, and so reduce the probability of a person taking weight control measures (Wardle, Haase, & Steptoe, 2006; Wetmore & Modkdad, 2012). This could potentially be treated by a training program which would improve discrimination of heavier bodies (the development of the "expertise effect" which reduce the JND for higher BMI bodies).

This study uses CGI 3D modelling to produce photorealistic bodies with a biometrically accurate simulation of weight change. These techniques are in increasing use as the modelling software becomes more user friendly to non-computer scientists and at the same time produces more realistic results (e.g., Evans, Tovée, Boothroyd, & Drewett, 2013; Frederick, Hadji-Michael, Furnham, & Swami, 2010; Glauert, Rhodes, Byrne, Fink, & Grammer, 2009; Tovée, Edmonds, & Vuong, 2012). This allows the use of sets of bodies which hold constant such features as their skin texture, proportions and facial features. However, we would underline the importance of biometric validation of these images as the software used to generate these images does not produce simple linear changes in body size

and without careful calibration of the resultant bodies, they may not be biometrically accurate and therefore may introduce a whole new set of artefacts into body research.

In conclusion, our results suggest that body size over-estimation is not only much more variable than might have been appreciated previously, given the strong dependence on individual BMI, but it also shows striking differences between women with ANSD and controls. The accuracy of control participants can be explained by the combined effects of two perceptual functions, contraction bias and Weber's law, with a modulating effect of psychological concerns about body size. By comparison, as their BMI increases, the accuracy of estimation by women with ANSD seems to be predicted primarily by a rapid criterion shift which is to compensate for a rapid decline in sensitivity to body size change.

### References

- Beck, A.T., Ward, C.H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, 4, 561-571.
- Berkman, N.D., Lohr, K.N., & Bulik, C.M. (2007). Outcomes of eating disorders: a systematic review of the literature. *International Journal of Eating Disorders*, 40, 293-309. doi: 10.1002/eat.20369
- Boothroyd, L.G., Tovée, M.J., & Pollet, T.V. (2012). Visual diet versus associative learning as mechanisms of change in body size preferences. *PLoS One*, 7, e48691. doi: 10.1371/journal.pone.0048691
- Cash, T.F., & Deagle, E.A. (1997). The nature and extent of body-image disturbances in anorexia nervosa and bulimia nervosa: A meta-analysis. *International Journal of Eating Disorders*, 22, 107-126.



- 1  
2  
3  
4 Casper, R.C., Halmi, K.A., Goldberg, S.C., Eckert, E., & Davis, J. (1979). Disturbances in  
5  
6 body image estimation as related to other characteristics and outcome in anorexia  
7  
8 nervosa. *The British Journal of Psychiatry*, *134*, 60-66.  
9
- 10  
11  
12 Channon, S., & DeSilva, W. (1985). Psychological correlates of weight gain in patients with  
13  
14 anorexia nervosa. *Journal of Psychiatric Research*, *19*, 267-271.  
15  
16
- 17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Cornelissen, P.L., Toveé, M.J., & Bateson, M. (2009). Patterns of subcutaneous fat  
deposition and the relationship between body mass index and waist-to-hip ratio:  
Implications for models of physical attractiveness. *Journal of Theoretical Biology*,  
*256*, 343-350. doi:10.1016/j.jtbi.2008.09.041
- Cornelissen, P.L., Johns, A., & Toveé, M.J. (2013). Body size over-estimation in women with  
anorexia nervosa is not qualitatively different from female controls. *Body Image*, *10*,  
103-111. doi: 10.1016/j.bodyim.2012.09.003
- Durnin, J., & Womersley, J. (1974). Body fat assessed from total body density and its  
estimation from skinfold thickness: measurements on 481 men and women aged from  
16 to 72 years. *British Journal of Nutrition*, *32*, 77-97.
- Evans, C., & Dolan, B. (1993). Body Shape Questionnaire: derivation of shortened “alternate  
forms”. *International Journal of Eating Disorders*, *13*, 315-321.
- Evans, E.H., Toveé, M.J., Boothroyd, L.G., & Drewett, R.F. (2013). Body dissatisfaction and  
disordered eating attitudes in 7- to 11-year old girls: testing a sociocultural model.  
*Body Image*, *10*, 8-15. doi: 10.1016/j.bodyim.2012.10.001
- Fairburn, C.G., & Beglin, S.J. (1994). Assessment of eating disorders: Interview or  
self-report questionnaire? *International Journal of Eating Disorders*, *16*, 363-370.

- 1  
2  
3  
4 Fairburn, C.G., Cooper, Z., & Shafran, R. (2003). Cognitive behaviour therapy for eating  
5  
6 disorders: A “transdiagnostic” theory and treatment. *Behaviour Research and*  
7  
8 *Therapy, 41*, 509-528. doi:10.1016/S0005-7967(02)00088-8  
9
- 10  
11 Fairburn, C.G., & Harrison, P.J. (2003). Eating disorders. *Lancet, 361*, 407–16.  
12  
13 doi: 10.1016/S0140-6736(03)12378-1  
14  
15  
16
- 17 Fernández, F., Probst, M., Meerman, R., & Vandereycken, W. (1994). Body size estimation  
18  
19 and body dissatisfaction in eating disorder patients and normal controls. *International*  
20  
21 *Journal of Eating Disorders, 16*, 307-310.  
22  
23  
24
- 25 Frederick, D.A., Hadji-Michael., M, Furnham, A., & Swami, V. (2010). The influence of leg-  
26  
27 to-body ratio (LBR) on judgments of female physical attractiveness: assessments of  
28  
29 computer-generated images varying in LBR. *Body Image, 7*, 51-55.  
30  
31 doi:10.1016/j.bodyim.2009.09.001  
32  
33  
34
- 35 Garner, D.M., & Garfinkel, P.E. (1979). The eating attitudes test: an index of the symptoms  
36  
37 of anorexia nervosa. *Psychological Medicine, 9*, 273-279.  
38  
39
- 40 Gardner, R.M. (1996). Methodological issues in assessment of the perceptual component of  
41  
42 body image disturbance. *British Journal of Psychology, 87*, 327-337.  
43  
44  
45
- 46 Gardner, R.M., & Moncrieff, C. (1988). Body image distortion in anorexics as a non-sensory  
47  
48 phenomenon: A signal detection approach. *Journal of Clinical Psychology, 44*, 101-  
49  
50 107.  
51  
52  
53
- 54 Gardner, R.M., & Bokenkamp, E.D. (1996). The role of sensory and nonsensory factors in  
55  
56 body size estimations of eating disorder subjects. *Journal of Clinical Psychology, 52*,  
57  
58 3-15.  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 Gardner, R.M., Brown, D.L., & Boice, R. (2012). Using amazon's mechanical turk website to  
5  
6 measure accuracy of body size estimation and body dissatisfaction. *Body image*, 9,  
7  
8 532-534. doi:10.1016/j.bodyim.2012.06.006  
9
- 10  
11 Gescheider, G.A. (1997). *Psychophysics: The Fundamentals*. (3<sup>rd</sup> Ed.) Lawrence Erlbaum,  
12  
13 New Jersey.  
14  
15  
16
- 17 Glauert, R., Rhodes, G., Byrne, S., Fink, B., & Grammer, K. (2009). Body dissatisfaction and  
18  
19 the effects of perceptual exposure on body norms and ideals. *International Journal of*  
20  
21 *Eating Disorders*, 42, 443-452. doi: 10.1002/eat.20640  
22  
23  
24
- 25 Hasler, N., Stoll, C., Sunkel, M., Rosenhahn, B., & Seidel, H.P. (2009). A statistical model of  
26  
27 human pose and body shape. *Computer Graphics Forum*, 28, 337-346.  
28  
29 doi: 10.1111/j.1467-8659.2009.01373.x  
30  
31
- 32  
33 Kaye, W.H., Fudge, J.L., & Paulus, M. (2009). New insights into symptoms and neurocircuit  
34  
35 function of anorexia nervosa. *Nature Reviews Neuroscience*, 10, 573-584.  
36  
37 doi:10.1038/nrn2682.  
38  
39
- 40 Kuchler, F., & Variyam, JN (2003). Mistakes were made: misperception as a barrier to  
41  
42 reducing overweight. *International Journal of Obesity*, 27, 856–86.  
43  
44 doi:10.1038/sj.ijo.0802293  
45  
46  
47
- 48 Kuskowska-Wolk, A., & Rössner, S. (1989). The “true” prevalence of obesity: a comparison  
49  
50 of objective weight and height measures versus self-reported and calibrated data.  
51  
52 *Scandinavian Journal of Primary Health Care*, 7, 79-82.  
53  
54  
55
- 56 Leopold, D.A., O'Toole, A.J., Vetter, T., & Blanz, V. (2001). Prototype-referenced shape  
57  
58 encoding revealed by high-level aftereffects. *Nature Neuroscience*, 4, 89–94.  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 Maximova, K., McGrath, J.J., Barnett, T., O'Loughlin, J., Paradis, G., & Lambert, M. (2008).  
5  
6 Do you see what I see? Weight status misperception and exposure to obesity among  
7  
8 children and adolescents. *International Journal of Obesity*, *32*, 1008-1015. doi:  
9  
10 10.1038/ijo.2008.15.  
11  
12  
13  
14 McCabe, M.P., Ricciardelli, L.A., & Ridge, D. (2006). "Who thinks I need a perfect body?"  
15  
16 Perceptions and internal dialogue among adolescents about their bodies. *Sex Roles*,  
17  
18 *55*, 409-419. doi: 10.1007/s11199-006-9093-0  
19  
20  
21  
22 Meerman, R. (1983). Experimental investigation of disturbances in body image estimation in  
23  
24 anorexia nervosa patients, and ballet and gymnastics pupils. *International Journal of*  
25  
26 *Eating Disorders*, *2*, 91-100.  
27  
28  
29  
30 Mond, J.M., Hay, P., Rodgers, B., & Owen, C. (2006). Eating disorder examination  
31  
32 questionnaire (EDE-Q): norms for young adult women. *Behaviour Research and*  
33  
34 *Therapy*, *44*, 53-62. doi:10.1016/j.brat.2004.12.003  
35  
36  
37  
38 Mussap, A.J., McCabe, M.P., & Ricciardelli, L.A. (2008). Implications of accuracy,  
39  
40 sensitivity, and variability of body size estimations to disordered eating. *Body Image*,  
41  
42 *5*, 80-90. doi: 10.1016/j.bodyim.2007.07.003  
43  
44  
45  
46 Norris, M.L., Boydell, K.M., Pinhas, L., & Katzman, D.K. (2006). Ana and the internet: A  
47  
48 review of pro-anorexia websites. *International Journal of Eating Disorders*, *39*, 443-  
49  
50 447. doi: 10.1002/eat.20305  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 Pike, K.M. (1998). Long-term course of anorexia nervosa: response, relapse, remission, and  
5  
6 recovery. *Clinical Psychology Review*, 18, 447-475.  
7  
8  
9  
10 Poulton, E.C., (1989). *Bias in quantifying judgements*. Hove: Erlbaum.  
11  
12  
13 Probst, M., Vandereycken, W., & Van Coppenolle, H. (1997). Body-size estimation in eating  
14  
15 disorders using video distortion on a life-size screen. *Psychotherapy and*  
16  
17 *Psychosomatics*, 66, 87-91.  
18  
19  
20 Probst, M., Vandereycken, W., Van Coppenolle, H., & Pieters, G. (1998). Body size  
21  
22 estimation in anorexia nervosa patients: the significance of overestimation. *Journal of*  
23  
24 *Psychosomatic Research*, 44, 451-456.  
25  
26  
27  
28 Ransom, D.C., La Guardia, J.G., Woody, E.Z., & Boyd, J.L. (2010). Interpersonal  
29  
30 interactions on online forums addressing eating concerns. *International Journal of*  
31  
32 *Eating Disorders*, 43, 161-170. doi: 10.1002/eat.20629  
33  
34  
35  
36 Robinson, E., & Kirkham, T.C. (2013). Is he a healthy weight? Exposure to obesity changes  
37  
38 perception of the weight status of others. *International Journal of Obesity*, 38, 663-  
39  
40 667. doi:10.1038/ijo.2013.154  
41  
42  
43  
44 Rosenberg, M. (1965). *Society and the adolescent self-image*. Princeton, NJ: Princeton  
45  
46 University Press.  
47  
48  
49 Slade, P., & Russell, G. (1973). Awareness of body dimensions in anorexia nervosa: cross-  
50  
51 sectional and longitudinal studies. *Psychological Medicine*, 3, 188-199.  
52  
53  
54  
55 Stice, E., & Shaw, H.E. (2002). Role of body dissatisfaction in the onset and maintenance of  
56  
57 eating pathology: a synthesis of research findings. *Journal of Psychosomatic*  
58  
59 *Research*, 53, 985-993. doi: 10.1016/S0022-3999(02)00488-9  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Striegel-Moore, R.H., Franko, D.L., Thompson, D., Barton, B., Schreiber, G.B., & Daniels,  
5  
6 S.R. (2004). Changes in weight and body image over time in women with eating  
7  
8 disorders. *International Journal of Eating Disorders*, 36, 315-327.

9  
10 doi: 10.1002/eat.20053

11  
12  
13  
14 Swinburn, B.A., Sacks, G., Hall, K.D., McPherson, K., Finegood, D.T., Moodie, M.L., &  
15  
16 Gortmaker, S.L. (2011). The global obesity pandemic: shaped by global drivers and  
17  
18 local environments. *Lancet*, 378, 804–814. doi: 10.1016/S0140-6736(11)60813-1

19  
20  
21  
22 Tovée, M.J., Benson, P.J., Emery, J.L., Mason, S.M., & Cohen-Tovée, E.M. (2003).

23  
24 Measurement of body size and shape perception in eating-disordered and control  
25  
26 observers using body-shape software. *British Journal of Psychology*, 94, 501-516.

27  
28 doi: 10.1348/000712603322503060

29  
30  
31  
32 Tovée, M.J., Edmonds, L., & Vuong, Q.C. (2012). Categorical perception of human female  
33  
34 physical attractiveness and health. *Evolution and Human Behavior*, 33, 85-93. doi:  
35  
36 10.1016/j.evolhumbehav.2011.05.008.

37  
38  
39  
40 Tovée, M.J., Swami, V., Furnham, A., & Mangalparsad, R. (2006). Changing perceptions of  
41  
42 attractiveness as observers are exposed to a different culture. *Evolution and Human*  
43  
44 *Behavior* 2006, 27(6), 443-456. doi: 10.1016/j.evolhumbehav.2006.05.004

45  
46  
47 Treasure, J., Claudino, A.M., & Zucker, N. (2010). Eating disorders. *Lancet*, 375, 583–593.

48  
49 doi: [http://dx.doi.org/10.1016/S0140-6736\(09\)61748-7](http://dx.doi.org/10.1016/S0140-6736(09)61748-7)

50  
51  
52  
53 Truesdale, K., & Stevens, J. (2008). Do the obese know they are obese? *North Carolina*  
54  
55 *Medical Journal*, 69, 188–194.

56  
57  
58 Wagner, A., Aizenstein, H., Venkatraman, V., Fudge, J., May, J., Mazurkewicz, L., ... Kaye,  
59  
60 W.H. (2007). Altered reward processing in women recovered from anorexia nervosa.

1  
2  
3  
4 *American Journal of Psychiatry*, 164, 1842-1849. doi:  
5  
6 10.1176/appi.ajp.2007.07040575.  
7  
8

9  
10 Wardle, J., Haase, A.M., & Steptoe, A. (2006). Body image and weight control in young  
11  
12 adults: international comparisons in university students from 22 countries.  
13  
14 *International Journal of Obesity*, 30, 644–651. doi:10.1038/sj.ijo.0803050  
15  
16

17 Wells, J.C., Treleaven, P., & Cole, T.J. (2007). BMI compared with 3-dimensional body  
18  
19 shape: the UK national sizing survey. *The American Journal of Clinical Nutrition*, 85,  
20  
21 419-425. doi:10.1038/sj.ijo.0803727.  
22  
23

24  
25 Wetmore, C., & Modkdad, A.H. (2012). In denial: misperceptions of weight change among  
26  
27 adults in the United States. *Preventative Medicine*, 56, 93–100.  
28  
29 doi:10.1038/sj.ijo.0803727  
30  
31

32  
33 Winkler, C., & Rhodes, G. (2005). Perceptual adaptation affects attractiveness of female  
34  
35 bodies. *British Journal of Psychology*, 96, 141-154. doi: 10.1348/000712605X36343  
36  
37  
38  
39  
40  
41  
42  
43

#### 44 **Acknowledgements**

45  
46  
47 KKC was supported by a PhD studentship funded by Northumbria University; We  
48  
49 thank: B-Eat, NIWE for help with the recruitment of participants; Dr. Nick Neave for help  
50  
51 with implementing the study; Jonathan Prichard for help with biometric measurements; Dr  
52  
53 Vesa Kiviniemi for advice on statistical analysis.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Figure Captions

### Figure 1

(A) Schematic representation of the results from Cornelissen et al. (2013); (B) The pattern of body size estimation predicted by the contraction bias model in women with AN, or recovering from AN (i.e., in an eating disordered group with a wider BMI range); (C) The pattern of body size over estimation predicted by increasing psychological concerns, rather than contraction bias. See text for details.

### Figure 2

The top row shows three CGI images from our database of stimuli. The second row shows the body shape changes produced by applying the video distortion technique (VDT) to the central image in the first row. The third row shows the difference between the two methods of image manipulation. See text for details.

### Figure 3

(A) Shows the relationship between participants' BMI (x-axis) and their subjective estimate of body size (PSE) with the effects of PSYCH statistically controlled. (B) Shows the relationship between participants' BMI (x-axis) and fitted values of estimated body size (PSE) computed from the mixed model at  $\pm 1$  SD of the mean PSYCH value for each group. (C) The relationship between participants' BMI (x-axis) and their sensitivity at estimating body size (Log10 DL) with the effects of PSYCH statistically controlled. (D) A sketch plot to illustrate the relationship between the psychometric function for body size estimation and the calculation of PSE (i.e., the value of BMI at  $y$  and  $y'$  when percentage 'larger' responses = 50%) and DL (i.e., the difference in BMI between  $z$  &  $x$  and  $z'$  &  $x'$ ). These points correspond



1  
2  
3  
4 to values of BMI when percentage 'larger' responses equal 75% and 25%, respectively). Two  
5  
6 situations are illustrated: accurate and sensitive performance (black) versus over-estimating  
7  
8 and insensitive performance (white).  
9

**Table 1**

Means and standard deviations of the characteristics of the three ANSD subgroups.

	ANSD1 (n=12)		ANSD2 (n=20)		ANSD3 (n=10)		ANSD 1v2	ANSD 1v3	ANSD 2v3
	M	SD	M	SD	M	SD	P	P	p
Participant characteristics									
Age (years)	22.42	3.65	25.80	7.62	22.36	3.44	.78	.99	.81
BMI (weight/height <sup>2</sup> )	18.40	2.35	21.03	2.95	22.62	1.59	.12	<.01	.63
Age at onset (years)	17.42	3.92	15.55	2.01	16.45	1.97	.54	.99	.92
When treatment ended (years)	0.00	0.00	3.45	4.11	1.95	2.90	-	.23	.96
Duration of illness (years)	5.04	4.70	6.75	6.79	4.91	3.05	.99	.99	.99
Depression and self esteem									
RSE score	8.33	2.35	16.58	5.03	15.70	5.81	<.01	.02	.99
BDI score	31.91	9.49	15.74	8.10	15.00	10.46	<.01	.01	.99
Eating and body shape concern									
EAT score	39.25	15.70	21.10	13.27	28.80	15.51	.01	.72	.80
BSQ score	71.00	17.69	55.50	17.00	62.20	18.47	.16	.95	.98
EDEQ score	4.01	1.34	2.75	1.18	3.05	1.65	.09	.76	.99
EDEQ wc score	4.08	1.58	3.05	1.34	3.70	1.50	.40	.99	.92
EDEQ bs score	4.70	1.35	3.77	1.41	3.81	1.58	.50	.81	.99
Psychophysical performance									
PSE (weight/height <sup>2</sup> )	20.57	4.93	21.95	5.26	23.66	4.15	.99	.66	.98
OE i.e. PSE – BMI	2.17	3.25	1.10	2.83	1.04	3.03	.99	.77	.93
DL (weight/height <sup>2</sup> )	0.53	0.28	0.65	0.46	0.98	1.05	.97	.99	.99

Note. ANSD1: AN participants being treated at the time of testing. ANSD 2: AN participants no longer being treated; ANSD 3: atypical AN or Bulimia that switched to AN. BDI = Beck Depression Inventory. RSE = Rosenberg Self-Esteem Scale. EAT = Eating Attitudes Test. BSQ = Body Shape Questionnaire. EDEQ = Eating Disorder Examination Questionnaire global score. EDEQ wc = Eating Disorder Examination Questionnaire weight concern subscale. EDEQ bs = Eating Disorder Examination Questionnaire body shape concern subscale. PSE = Point of Subjective Equality. OE = Over Estimation. DL = Difference Limen.

**Table 2:** Means and standard deviations for the participant characteristics, separated according to whether they belong to the ANSD or the healthy control group.

	ANSD (n=42)		Control(n=100)		p
	M	SD	M	SD	
Participant characteristics					
Age (years)	23.80	6.05	24.02	8.91	.99
BMI (weight/height <sup>2</sup> )	20.89	2.84	24.01	5.03	<.01
Percentage body fat	25.56	6.41	31.63	5.04	<.01
Depression and self esteem					
BDI	20.00	11.37	9.09	6.08	<.01
RSE	14.00	5.96	18.56	4.79	<.01
Eating and body shape concern					
EAT	27.38	15.55	11.07	9.08	<.01
BSQ	61.10	18.63	50.08	17.33	<.01
EDEQ	3.12	1.42	2.01	1.21	<.01
EDEQ wc	3.50	1.48	2.21	1.44	<.01
EDEQ bs	4.04	1.46	2.97	1.50	<.01
Psychophysical performance					
PSE (weight/height <sup>2</sup> )	22.21	4.95	23.80	5.13	.36
OE (i.e. PSE – BMI)	1.41	3.04	-0.14	2.85	.04
DL (weight/height <sup>2</sup> )	0.72	0.66	0.99	0.81	.32

Note. BDI = Beck Depression Inventory. RSE = Rosenberg Self-Esteem Scale. EAT = Eating Attitudes Test. BSQ = Body Shape Questionnaire. EDEQ = Eating Disorder Examination Questionnaire global score. EDEQ wc = Eating Disorder Examination Questionnaire weight concern subscale. EDEQ bs = Eating Disorder Examination Questionnaire body shape concern subscale. PSE = Point of Subjective Equality. OE = Over Estimation. DL = Difference Limen.

Figure-1  
[Click here to download high resolution image](#)




Figure-2  
[Click here to download high resolution image](#)

BMI ~13

BMI ~17

BMI ~21

CGI



VDT



Difference  
between  
CGI  
and VDT



 VDT  
 CGI



Figure-3  
[Click here to download high resolution image](#)

