

Northumbria Research Link

Citation: Jones, Emma, Sünram-Lea, Sandra and Wesnes, Keith (2012) Acute ingestion of different macronutrients differentially enhances aspects of memory and attention in healthy young adults. *Biological Psychology*, 89 (2). pp. 477-486. ISSN 0301-0511

Published by: UNSPECIFIED

URL:

This version was downloaded from Northumbria Research Link: <http://northumbria-test.eprints-hosting.org/id/eprint/47705/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



UniversityLibrary



Northumbria
University
NEWCASTLE

Northumbria Research Link

Citation: Jones, Emma, Sünram-Lea, Sandra and Wesnes, Keith (2012) Acute ingestion of different macronutrients differentially enhances aspects of memory and attention in healthy young adults. *Biological Psychology*, 89 (2). pp. 477-486. ISSN 0301-0511

Published by: Elsevier

URL: <http://dx.doi.org/10.1016/j.biopsycho.2011.12.017>
<<http://dx.doi.org/10.1016/j.biopsycho.2011.12.017>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/5524/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



Manuscript Number: BIOPSY-D-10-00384R2

Title: Acute ingestion of different macronutrients differentially enhances aspects of memory and attention in healthy young adults

Article Type: Research Paper

Keywords: macronutrients; cognitive function; memory; attention; glucose; protein; fat

Corresponding Author: Dr Emma Kathleen Jones, Ph.D.

Corresponding Author's Institution: PaCT Research centre

First Author: Emma Kathleen Jones, Ph.D.

Order of Authors: Emma Kathleen Jones, Ph.D.; Sandra I Sünram-Lea, Ph.D.; Keith A Wesnes, Ph.D. FSS
CPsychol FBPsS

Abstract: The role of carbohydrates on mood and cognition is fairly well established, however research examining the behavioural effects of the other macronutrients is limited. The current study compared the effects of a 25g glucose drink to energetically-matched protein and fat drinks and an inert placebo. Following a blind, placebo-controlled, randomized crossover design, 18 healthy young adults consumed drinks containing fat, glucose, protein and placebo. Cognitive performance was examined at baseline and again 15- and 60-minutes post drink. Mood was assessed at baseline and then 10-, 35- and 80-minutes post drink. Attention and speed were enhanced 15-minutes following fat or glucose ingestion and working memory was enhanced 15-minutes following protein ingestion. Sixty minutes post drink memory enhancements were observed after protein and memory impairment was observed following glucose. All drinks increased ratings of alertness. The findings suggest that macronutrients: i) have different windows of opportunity for effects ii) target different cognitive domains.

Dear Professor Ring,

Thank you for inviting us to respond to the final reviewer comment and resubmit our paper to *Biological Psychology*. Below we have addressed and outlined in detail our response to the comment made by reviewer 2, which we had not sufficiently addressed previously. We hope you find our comments and amendments acceptable and our paper now worthy of publication.

Best wishes

Emma Jones

Response to Review:

When selecting the current design we acknowledged that both between-subjects and repeated measures have their own problems and it is necessary to identify and attempt to reduce these as much as possible. We selected a crossover design, the particular strength of which is that the interventions under investigation are evaluated within the same participant and so eliminates between-subject variability (Maclure, 1991). However, we do agree that the design has certain weaknesses, including carry-over of effect of treatments across study periods, which could potentially distort the results (Cleophas, 1990; Wallenstein & Fisher, 1977) and observed treatment effects will depend upon the order in which they were received.

Some have argued against consistent testing for carryover effects of interventions across periods as carry-over effects are rare and statistical manipulation after the fact cannot address the impact of a carry-over effect (Senn, D'Angelo & Potvin, 2004). It has been argued that tests for carry-over are generally underpowered even with an appreciable carry-over effect (Senn, 1988). Another complicating matter when assessing order effects is that effects may interact with participant, and this is difficult to assess. Consequently any effect of order or interaction effects may not detect the order effect.

However what is of particular relevance in the current study is that statistical analysis was carried out on change from baseline levels, which in itself controls for potential carry over effects. The treatment we used is not an endogenous entity nor has administration of fat, protein and glucose any long lasting effects. Moreover, we employed a 5-7 day washout between study days and treatment order was randomised. In addition, to reduce potential effect of familiarity with tests, the current study employed four practice sessions prior to the start of the study.

We have however followed the recommendation of reviewer 2 and incorporated order in the ANOVA model and the results showed no significant effects of drink order on any of the measures and only one significant three-way interaction (drink*time*drink order) which was on quality of working memory [$F(3,36)=3.44$, $p=0.013$]. Given the number of "orders" it is unclear what this means. The drink*time interaction is also

significant (as already reported) and this has been explored further in the paper. We have highlighted this issue in the discussion section of the paper (first para. Pg 20).

Consequently, although we do appreciate the concerns of reviewer 2, we are confident that the design of the study (practice session, randomisation, pre-and post treatment assessment) minimised order effects and is overall preferable to the potential confounding variables that present with between participant designs.

References

Maclure M: The case-crossover design: a method for studying transient effects on the risk of acute events. *Am J Epidemiol.* 1991, 133(2):144-153.

Cleophas TJ: A simple method for the estimation of interaction bias in crossover studies. *J Clin Pharmacol* 1990, 30:1036-1040

Wallenstein S, Fisher AC: The analysis of the two-period repeated measurements crossover design with application to clinical trials. *Biometrics* 1977, 33:261-269.

Senn SJ, D'Angelo G, Potvin D: Carry-over in cross-over trials in bioequivalence: theoretical concerns and empirical evidence. *Pharmaceutical Statistics* 2004, 3:13-142.

Senn SJ: Cross-over trials, carry-over effects and the art of self-delusion. *Stat Med* 1988, 7:1099-101

Dear Reviewer,

Thank you for your helpful comments and suggestions. I apologise that we failed to fully address your concerns in our previous response and hope we have dealt with the issue to your satisfaction below. We hope that our paper is now worthy of publication.

Best wishes

Emma Jones

Response to Reviewer:

When selecting the current design we acknowledged that both between-subjects and repeated measures have their own problems and it is necessary to identify and attempt to reduce these as much as possible. We selected a crossover design, the particular strength of which is that the interventions under investigation are evaluated within the same participant and so eliminates between-subject variability (Maclure, 1991). However, we do agree that the design has certain weaknesses, including carry-over of effect of treatments across study periods, which could potentially distort the results (Cleophas, 1990; Wallenstein & Fisher, 1977) and observed treatment effects will depend upon the order in which they were received.

Some have argued against consistent testing for carryover effects of interventions across periods as carry-over effects are rare and statistical manipulation after the fact cannot address the impact of a carry-over effect (Senn, D'Angelo & Potvin, 2004). It has been argued that tests for carry-over are generally underpowered even with an appreciable carry-over effect (Senn, 1988). Another complicating matter when assessing order effects is that effects may interact with participant, and this is difficult to assess. Consequently any effect of order or interaction effects may not detect the order effect.

However what is of particular relevance in the current study is that statistical analysis was carried out on change from baseline levels, which in itself controls for potential carry over effects. The treatment we used is not an endogenous entity nor has administration of fat, protein and glucose any long lasting effects. Moreover, we employed a 5-7 day washout between study days and treatment order was randomised. In addition, to reduce potential effect of familiarity with tests, the current study employed four practice sessions prior to the start of the study.

We have however followed the recommendation of reviewer 2 and incorporated order in the ANOVA model and the results showed no significant effects of drink order on any of the measures and only one significant three-way interaction (drink*time*drink order) which was on quality of working memory [$F(3,36)=3.44$, $p=0.013$]. Given the number of "orders" it is unclear what this means. The drink*time interaction is also significant (as already reported) and this has been explored further in the paper. We have highlighted this issue in the discussion section of the paper (first para. Pg 20).

Consequently, although we do appreciate the concerns of reviewer 2, we are confident that the design of the study (practice session, randomisation, pre-and post treatment assessment) minimised order effects and is overall preferable to the potential confounding variables that present with between participant designs.

References

Maclure M: The case-crossover design: a method for studying transient effects on the risk of acute events. *Am J Epidemiol.* 1991, 133(2):144-153.

Cleophas TJ: A simple method for the estimation of interaction bias in crossover studies. *J Clin Pharmacol* 1990, 30:1036-1040

Wallenstein S, Fisher AC: The analysis of the two-period repeated measurements crossover design with application to clinical trials. *Biometrics* 1977, 33:261-269.

Senn SJ, D'Angelo G, Potvin D: Carry-over in cross-over trials in bioequivalence: theoretical concerns and empirical evidence. *Pharmaceutical Statistics* 2004, 3:13-142.

Senn SJ: Cross-over trials, carry-over effects and the art of self-delusion. *Stat Med* 1988, 7:1099-101

*Highlights

- Attention and speed of processing were enhanced 15-minutes following fat or glucose ingestion
- Working memory was enhanced 15-minutes following protein ingestion
- Sixty minutes post drink memory enhancements were observed after protein and memory impairment was observed following glucose
- All drinks (including placebo) increased ratings of alertness immediately post drink

Running head: ACUTE MACRONUTRIENT INGESTION ENHANCES COGNITION

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acute ingestion of different macronutrients differentially enhances aspects of memory and
attention in healthy young adults

Emma K. Jones^{a1}, Sandra I Sünram-Lea^a and Keith A. Wesnes^b

- a. Department of Psychology, Fylde College, University of Lancaster, Lancaster LA1
4YF, UK. emma2.jones@northumbria.ac.uk
- b. United BioSource Corporation, Gatehampton Road, Goring on Themes RG8 0EN,
UK

Corresponding author: Emma Jones, Psychology and Communication Technology
Research Centre, Department of Psychology, Northumbria University, Newcastle upon
Tyne NE1 8ST, UK.
Email: emma2.jones@northumbria.ac.uk
Tel: 0191 227 3723

¹ Present address: Psychology and Communication Technology Research Centre, Department
of Psychology, Northumbria University, Newcastle upon Tyne NE1 8ST, UK.

Abstract

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The role of carbohydrates on mood and cognition is fairly well established, however research examining the behavioural effects of the other macronutrients is limited. The current study compared the effects of a 25g glucose drink to energetically-matched protein and fat drinks and an inert placebo. Following a blind, placebo-controlled, randomized crossover design, 18 healthy young adults consumed drinks containing fat, glucose, protein and placebo. Cognitive performance was examined at baseline and again 15- and 60-minutes post drink. Mood was assessed at baseline and then 10-, 35- and 80-minutes post drink. Attention and speed were enhanced 15-minutes following fat or glucose ingestion and working memory was enhanced 15-minutes following protein ingestion. Sixty minutes post drink memory enhancements were observed after protein and memory impairment was observed following glucose. All drinks increased ratings of alertness. The findings suggest that macronutrients: i) have different windows of opportunity for effects ii) target different cognitive domains.

Introduction

1
2
3 The effects of nutrition on brain and behaviour and more specifically the
4
5 cognitive effects of foods, food components and nutritional interventions are very
6
7 much on the public agenda. More specifically, it has been demonstrated that acute
8
9 administration of glucose (a simple carbohydrate), can facilitate verbal declarative
10
11 memory in healthy young adults and adolescents (e.g. Foster, Lidder & Sünram, 1998;
12
13 Smith & Foster, 2008; Smith Hii, Foster & van Eekelen, 2009; Sünram-Lea, Foster,
14
15 Durlach & Perez, 2001; 2002a; 2002b) and older populations (e.g. Craft, Murphy &
16
17 Wemstrom, 1994; Kaplan, Greenwood, Winocur & Wolever, 2001; Messier, Gagnon
18
19 & Knott, 1997; Riby, Meikle & Glover, 2004). In addition, administration of a
20
21 glucose drink has been shown to improve working memory performance in healthy,
22
23 young adults (e.g. Scholey, Harper & Kennedy, 2001); to ameliorate impairment of a
24
25 secondary, psychomotor task during a divided attention (encoding plus psychomotor)
26
27 paradigm (Scholey, Sünram-Lea, Greer, Elliott, & Kennedy, 2009) and to enhance
28
29 prospective memory (Riby, Laws, McLaughlin & Murray, 2011). Hoyland, Lawton
30
31 and Dye (2008) conducted a comprehensive review of the literature and concluded
32
33 that the most robust glucose-mediated enhancement has been demonstrated on
34
35 memory although there are numerous examples of glucose facilitation of other
36
37 cognitive tasks (see Hoyland et al., 2008; Messier, 2004; Riby, 2004 for reviews).
38
39
40
41
42
43
44
45

46 Glucose index (GI) is the rate at which an ingested substance increases and
47
48 maintains blood glucose levels. Pure glucose has a high GI as it increases levels
49
50 quickly with a fast return to baseline, whereas foodstuff with lower GI ratings tend to
51
52 elicit a slower rise, smaller peak and are slower to return to baseline. Glycaemic Load
53
54 (GL) is another way of describing response to a carbohydrate and takes into account
55
56 both GI and the quantity of food. It has been found that foods with different GIs and
57
58
59
60
61
62
63
64
65

1 GLs can differentially influence cognitive function. For example, some authors report
2 that cognitive performance benefits can be more readily observed following
3
4 consumption of low GI foods compared to high GI foods (e.g. Benton et al., 2003).
5
6 More specifically, researchers have demonstrated beneficial effects of low GI
7
8 breakfasts compared to high GI breakfasts for children (e.g. Ingwersen, Defeyter,
9
10 Kennedy, Wesnes & Scholey, 2007; Mahoney, Taylor, Kanarek & Samuel, 2005;
11
12 Wesnes, Pincock, Richardson, Helm & Hails, 2003). Ingwersen et al. (2007) found
13
14 that following a low GI breakfast cognitive benefits were observed for 2-hours post
15
16 ingestion whereas performance following the low GI breakfast declined over this time
17
18 period. A review by Gilsenan and colleagues (Gilsenan, Bruin & Dye, 2009)
19
20 concluded that evidence for effects of different GLs over relatively short periods of
21
22 time (from between 100-390 minutes) is inconsistent. However, in general it appears
23
24 that high GI or GL foods (including glucose drinks) appear to have short term
25
26 cognitive benefit, whereas over a longer time frame foods which allow a more
27
28 sustained energy supply are more beneficial (also see e.g. Kaplan, Greenwood,
29
30 Winocur & Wolver, 2000).
31
32
33
34
35
36
37
38

39 To date, a large body of literature reports on the influence of glucose ingestion
40
41 on cognitive function whereas considerably fewer studies have examined the
42
43 influence of protein or fat ingestion and/or compared the effects of different
44
45 macronutrients on cognitive performance. Kaplan et al., (2001) compared the effects
46
47 of protein, glucose and fat ingestion in a sample of older participants (61–79 years)
48
49 and found that immediately after ingestion all three macronutrients improved memory
50
51 performance compared to placebo, whereas 60-minutes post-ingestion memory
52
53 improvements were only observed following a glucose drink. In addition, fat
54
55 administration improved attention 60 minutes post ingestion and protein led to a
56
57
58
59
60
61
62
63
64
65

1 reduced rate of forgetting when assessed 15 min after consumption. Fischer,
2 Colombani, Langhans and Wenk (2001) administered carbohydrate-, protein- and fat-
3 containing meals to healthy, young adults and tested cognitive performance at various
4 time-points over the course of 180-minutes. Their data indicated that different
5 macronutrients influence cognitive performance in a different manner, with the best
6 performance usually observed after fat ingestion. More specifically, faster reaction
7 times, improved short-term memory and improved attention were observed at all time
8 points (60-, 120- and 180 minutes) following fat ingestion. In a placebo-controlled
9 study Jones, Sünram-Lea and Threadgold (2005) compared the effects of glucose and
10 protein administration and observed glucose-mediated enhancement of reaction times
11 following glucose administration. However, this study failed to replicate the
12 immediate beneficial effects of macronutrients observed by Kaplan et al. or the well-
13 documented glucose facilitation of memory. Overall, findings to date tentatively
14 suggest that macronutrients may differentially influence cognition and mood and that
15 the effects are time-dependent (e.g. Fischer et al., 2001; Kaplan et al., 2001).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 In addition to cognitive facilitation, previous research has also demonstrated
37 complex macronutrient-specific effects on mood. For example, Fischer et al. (2001)
38 observed a reduction in depression scores on the Profile of Mood Scale (POMS)
39 following carbohydrate ingestion compared to protein. Conversely, Gibson et al.
40 (1999) found that 120-minutes post ingestion, a meal high in protein (and low in
41 carbohydrate) led to increased positive affect on the PANAS mood scale compared to
42 a meal low in protein (and high in carbohydrate). In our own laboratory we have
43 previously observed an increase in negative affect following protein ingestion (Jones
44 et al., 2005). These findings appear to be contradictory but methodological differences
45 between studies may account for these discrepancies. For example, Fischer et al.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 (2001) administered pure macronutrients whereas Gibson et al. (1999) administered
2 macronutrient combinations, which preclude a clear conclusion as to whether the
3 mood effects are due to specific macronutrients or interaction effects of multiple
4 macronutrients. In addition, the use of different mood rating scales further impedes
5 direct comparison.
6
7
8
9
10

11 It is evident from this brief review of the literature that the effects of glucose
12 on cognitive functioning and mood are relatively well described whereas those of
13 protein and fat still remain to be explored in detail. It is also important to note that
14 although investigating the effects of combined administration of different
15 macronutrients (for example Benton & Sargent, 1992) is important to for our
16 understanding of macronutrient interaction they do not inform us about the
17 contributions of individual macronutrients and their potential underlying mechanisms.
18 It is important to establish whether different macronutrients influence cognitive
19 performance via a specific mechanism or whether they exert their influence via a
20 shared, generalised mechanism. Kaplan et al. (2001) found evidence to indicate a
21 generalised effect of macronutrients on cognitive function in older adults (e.g. effects
22 15 and 60 minutes post ingestion - earlier than would be expected if metabolite-
23 mediated) suggesting a pre-digestive influence possibly due to the release of gut
24 hormones such as cholecystokinin (CCK), gastrin-releasing peptide or amylin which
25 have all been demonstrated to influence memory in animals (Flood and Morley, 1989;
26 1992; Flood, Smith & Morley, 1987; Morley, Flood, Silver & Kaiser, 1994). Whether
27 such effects have nutrient and/ or domain specificity is yet to be examined. There is
28 now growing interest in the potential of specific foods to influence mood and
29 cognitive performance. The main components of our diet that can be readily
30 manipulated are the macronutrients, glucose (carbohydrates), protein and fat.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Therefore, the aim of the current research was to further explore the effects of acute
2 macronutrient ingestion on cognitive performance and mood in healthy young adults.
3
4 Specifically, the effects of acute fat, protein, glucose and placebo ingestion on a range
5
6 of cognitive tests and mood scales were examined.
7

8 9 10 Methods and Materials

11 12 Power calculation

13
14 A medium overall effect size was previously found in a meta-analytic review
15
16 of the glucose facilitation effect ($d = 0.56$; Riby, 2004). An a priori power calculation
17
18 using G-power (Erdfelder, Faul, & Buchner, 1996) revealed that for a medium effect
19
20 size, with alpha set to 0.05 (two-tailed) a sample size of 16 would be required for 95%
21
22 power. The effect size of fat and protein is as yet unclear due to the relatively few
23
24 numbers of studies carried out in this area, however Kaplan et al. (2001) observed
25
26 effects using 22 participants.
27
28
29
30

31 32 Participants

33
34
35 Eighteen healthy young male and female participants (5 males, 13 females)
36
37 with a mean BMI of 21.1 kg/m^2 took part in this study. Ages ranged from 18-37 years
38
39 (mean age = 19 years). Participants were excluded from the study on the basis of several
40
41 criteria. Information regarding these criteria was gathered using a confidential medical
42
43 questionnaire which was completed before signing the consent form. Exclusion criteria
44
45 included i) history of neurological and/or psychiatric illness, ii) Diabetes Mellitus, iii)
46
47 BMI ≥ 25 , iv) intolerance or allergic reaction to substances that contain phenylalanine.
48
49
50 Participants were recruited from the University of Lancaster 1st year cohort and
51
52 received 30 pounds sterling and 4 course credits for taking part in the experiment. The
53
54 study was approved by the Ethics Committee of the Department of Psychology,
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 guidelines. Written informed consent was obtained from each participant prior to
2 participation.
3

4 Design

5
6
7 Following a blind, placebo controlled, balanced, cross-over, repeated measures
8 design, participants were administered 40g protein in solution, 16g fat emulsion, 40g
9 glucose solution and an inert placebo (matched for volume, sweetness and flavour)
10 over four study days, with a 5-7 day washout period between treatments. Treatment
11 order was randomly assigned. Treatment order was randomised and counterbalanced
12 using a Latin Square.
13
14
15
16
17
18
19
20
21

22 Treatments

23
24
25 Three isoenergetic (145 Kcal) and isovolumic (300ml) drinks and an inert
26 placebo were administered. All drinks were matched for volume, sweetness and
27 flavour and administered in opaque cups, covered by lids and ingested through a
28 straw. Drinks were flavoured with lemon juice in order to improve palatability and
29 participants' compliance. Drinks were prepared in the laboratory and refrigerated
30 prior to testing. The composition of test drinks is shown in Table 1.
31
32
33
34
35
36
37
38
39

40 <Table 1. here>
41
42

43 Blood Glucose Measurement

44
45
46 Blood glucose readings were obtained using the ExacTech blood glucose
47 monitoring equipment (supplied by MediSense Britain Ltd, 16/17 The Courtyard,
48 Gorsey Lane, Coleshill, Birmingham B46 1JA), following the manufacturers
49 recommended procedure.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Measuring Cognition and Mood

1
2
3 A tailored version of the CDR System (www.unitedbiosource.com) was used
4
5 to assess participants' cognitive performance and mood. The battery was administered
6
7 on PCs and responses were made via a two-button (yes and no) response box.
8

9 Responses for the Visual Analogue Scales (VAS) were made by mouse click.

10
11
12 Completion of the whole battery took around 20-minutes and tasks were presented in
13
14 following order:

15
16
17 Word presentation. – A list of 15 words matched for frequency, concreteness
18
19 and imagery was presented on the monitor at the rate of one every two seconds for
20
21 participants to remember. During encoding, participants were required to perform two
22
23 complex hand-movement sequences (Sünram-Lea et al., 2001). Each sequence was
24
25 performed using both hands and contained three movements: fist – chop - slap and
26
27 back-slap – chop – fist. Participants were told to alternate the sequence every fifth
28
29 word and they were not informed when to change, only that they had to keep track of
30
31 this themselves. Hand-movements were performed continually during word
32
33 presentation.
34
35
36
37
38

39 Immediate word recall. - Immediately after the words had been presented
40
41 participants were given 60-seconds to write down as many words as they could from
42
43 the list they had just seen. Participants' responses were marked according to total
44
45 number of errors, intrusions and percentage of words recalled correctly (accuracy).
46
47
48

49 Picture presentation. – Twenty photographs of objects were individually,
50
51 displayed in the centre of the screen at a rate of one every three seconds. Each picture
52
53 was displayed for one second. Participants were required to remember the pictures.
54
55

56 Simple reaction time. – The word 'yes' was presented repeatedly in the centre
57
58 of the screen with inter trial intervals varying randomly between 1 and 3.5 seconds.
59
60
61
62
63
64
65

1 Participants were required to respond by pressing the ‘yes’ button on their response
2 box as quickly as possible, whenever the word appeared. Reaction times were
3
4 recorded in milliseconds.
5

6
7 Digit vigilance. – A single target digit was randomly selected and
8
9 continuously displayed on the right side of the screen. In the centre a series of rapidly
10
11 changing digits was displayed at the rate of 150 digits per minute. Participants were
12
13 required to press the ‘yes’ button as quickly as possible, whenever the digit in the
14
15 centre matched the target digit. The task lasted for three minutes Reaction times
16
17 (milliseconds), percentage accuracy and number of false alarms were recorded.
18
19
20

21
22 Choice reaction time. - The target words ‘yes’ and ‘no’ were repeatedly,
23
24 randomly displayed individually in the centre of the screen. The inter-trial intervals
25
26 varied randomly between 1 second and 3.5 seconds. Participants were instructed to
27
28 respond by pressing the appropriate button on their response box as quickly and
29
30 accurately as possible. Reaction times (milliseconds) and percentage accuracy were
31
32 recorded.
33
34
35

36
37 Spatial working memory. - A picture of a house was displayed on the screen
38
39 with nine evenly distributed windows. Four of the windows were lit up in the original
40
41 picture and participants were asked to remember the position of these windows.
42
43 Following this, the house was presented again, repeatedly but each time only one
44
45 window was lit up. Participants were required to answer whether the window was lit
46
47 up or not in the original house by pressing the appropriate button on their answer box.
48
49 Percentage accuracy of identifying novel stimuli (distractors) and target stimuli were
50
51 recorded in addition to reaction times (milliseconds) to distractors and targets and
52
53 overall reaction times. Sensitivity Index (SI) was calculated by combining an
54
55 individual’s ability to discriminate targets and their ability to discriminate distractors.
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

SI ranges between +1 and -1 whereby +1 indicates perfect performance, zero indicates chance performance, and a negative score indicates performance which is worse than chance.

Numeric working memory. - A series of five digits were displayed individually on the screen for participants to remember. These were followed by 30 probe digits to which participants were required to respond using their answer box, indicating whether or not each probe digit had been in the original sequence. Percentage accuracy and reaction times for both distractors and targets were recorded in addition to overall reaction times and SI.

Delayed word recall. - Participants were given 60-seconds to write as many words as they could from the list they had seen at the beginning of the battery. Participant's responses were marked according to total number of errors, intrusions and percentage of words recalled correctly (accuracy).

Delayed word recognition. - The 15 original words and 15 distractor words were presented individually in a randomised order. Participants were asked to indicate whether each word had been in the original list or not by responding 'yes' or 'no' on their response box. Percentage accuracy and reaction times for both distractors and targets were recorded in addition to overall reaction times and SI.

Picture recognition. - The 20 original pictures and 20 distractor pictures were presented, individually in a randomised order. Participants were asked to indicate whether each word had been in the original list or not by responding 'yes' or 'no' on their response box. Percentage accuracy and reaction times for both distractors and targets were recorded in addition to overall reaction times and SI.

Subjective Mood

1 The Bond and Lader visual analogue scales (VAS) were used to assess
2 subjective mood (Bond & Lader, 1974). Sixteen VAS were presented on the screen
3 immediately after the cognitive tests. Participants used the mouse to position an arrow
4 at the point on the scale that represented their feelings at that moment. The 16 scales
5 were combined as recommended by Bond and Lader (1974) to form three mood
6 factors: 'alertness', 'calmness' and 'contentment'.
7
8
9
10
11
12
13

14 Cognitive Outcome Measures

15
16 Scores from individual measures were combined to form seven secondary
17 outcome measures ('power of attention', continuity of attention', 'quality of working
18 memory', 'quality of episodic secondary memory', 'quality of memory', 'speed of
19 memory' and 'combined speed') derived from factor analysis of the Cognitive Drug
20 Research computerised test battery (Wesnes, Ward, Ayre, & Pincock, 1999; Wesnes,
21 Ward, McGinty & Petrini, 2000), and previously used (e.g. Wesnes et al., 1997; 1999;
22 2000; Kennedy, Scholey, & Wesnes, 2001; 2002; Kennedy, Scholey, Tildsley, Perry
23 & Wesnes, 2002; Sünram-Lea, Birchall, Wesnes & Petrini, 2004). See Figure 1.
24
25
26
27
28
29
30
31
32
33

34
35
36 Power of attention *factor* (also referred to as '*speed of attention*'): Derived by
37 combining reaction times of three attention tasks: simple reaction time, choice
38 reaction time, and digit vigilance (units are summed milliseconds for the three
39 tasks).
40
41
42
43
44

45
46 Continuity of attention *factor* (also referred to as '*accuracy of attention*'):
47 Derived by calculating the combined percentage accuracy across choice reaction
48 time and digit vigilance tasks (with adjustment for false alarms on the latter
49 test). 100 percent accuracy across the two tasks would result in a maximum
50 score of 95.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Quality of working memory: Derived by combining SI scores from the two
2 working memory tests: spatial working memory and numeric working memory.

3
4
5 Range from -2 to +2. Perfect performance on both tasks result in a maximum
6 score of +2.

7
8
9 Quality of episodic secondary memory: Derived by calculating the combined
10 percentage accuracy scores (adjusted for proportion of novel and new stimuli
11 where appropriate) from all secondary memory tests: word recognition, picture
12 recognition, immediate word recall, delayed word recall (with adjustment to the
13 total percentage correct for errors and intrusions on the latter two tasks). One
14 hundred percent accuracy across the four tasks would result in a maximum score
15 of 400.

16
17
18 Quality of memory factor: Derived by calculating the combined percentage
19 accuracy scores (adjusted for proportion of novel and new stimuli where
20 appropriate) of all working memory tests and secondary memory tests: spatial
21 working memory, numeric working memory, word recognition, picture
22 recognition, immediate word recall, delayed word recall (with adjustment to the
23 total percentage correct for errors and intrusions on the latter two tasks). One
24 hundred percent accuracy across the six tasks would generate a maximum score
25 of 600.

26
27
28 Speed of memory factor: Derived by combining reaction times of the numeric
29 working memory task, spatial memory task, delayed word recognition and
30 delayed picture recognition task (units are summed milliseconds for the four
31 tasks).

32
33
34 Combined speed: Derived by combining the two speed outcome factors: 'speed
35 of memory' and 'power of attention'.

36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

<Figure 1 goes here>

Procedure

Each participant attended four 20-minute practice sessions in order to familiarise them with the cognitive test battery. Upon arrival at the laboratory for the first practice session, participants gave informed consent and demographic information. They were given complete instructions for each task including the secondary hand movement task. No treatments were administered during the practice sessions and performance data from these sessions was not included in the analysis.

Once the practice sessions had been successfully completed there were four experimental sessions. All followed the same procedure. On arrival at their first session participants were randomly allocated to a treatment regime which counterbalanced the order of drinks across the study days. Sessions were separated by a 5-7 day wash-out period and they were conducted in the mornings following a 12-hour, over-night fast. In addition, participants were instructed to refrain from nicotine, alcohol and stimulants for 12-hours prior to each session. Sessions were 1 hour and 45 minutes long and started at either 9am or 11am.

On entering the laboratory participants were asked to report if they had complied to the 12-hour fast. The first blood glucose measure (T0) was taken and the cognitive test battery was administered (pre-treatment assessment) followed by drink administration. 10-minutes were allowed for drink consumption and immediately following consumption the visual analogue scales were administered to assess any immediate effects of drink ingestion on mood. The second blood glucose measure was taken 12-minutes following drink ingestion (T12). 15-minutes following drink administration the cognitive test battery and mood scales were administered again.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Upon completion of the test battery and mood scales the third blood glucose reading was taken (T37). Participants were then allowed to engage in silent reading of their own choice for 20-25 minutes, after which the fourth blood glucose reading was taken (T55). Sixty minutes post-drink, the cognitive battery was, again, administered, followed by the last blood glucose measure (T82). This time-scale was employed in this study as glucose facilitation appears to be optimal when testing starts 15 to 20 minutes post ingestion (e.g. Foster et al., 1998; Owens & Benton, 1994; Sünram-Lea et al., 2001; 2002a; 2002b; 2004), whereas beneficial effects of fat and protein ingestion have been observed following longer delays (Kaplan et al., 2001; Fischer et al., 2001)

Statistical Analysis

Blood Glucose Levels

Blood glucose levels (mmol/litre) were analysed using a two-way (4*5) repeated measures ANOVA (drink: fat, protein, glucose and placebo and time: T0, T12, T37, T55 and T82). Significant main effects and interactions were analysed using the Bonferroni post hoc test.

Cognitive Data and Visual Analogue Scales

Scores on the cognitive outcome measures and the three factors derived from the visual analogue scales were analysed as 'change from baseline'. Comparisons of all drinks were made using repeated measures ANOVAs. To further explore main effects and interactions, planned comparisons of each treatment drink with placebo were made using t-tests with MSE from omnibus ANOVA as an error term. Post hoc comparisons were made using Bonferroni tests. For all cognitive outcome factors the ANOVAs were 4*2 (drink: fat, protein, glucose and aspartame and assessment: 15-

1 minutes post and 60-minutes post). For the visual analogue scales the levels of
2 assessment were: 10-minutes post, 35-minutes post and 80-minutes post. In order to
3
4 minimise the risk of type 1 errors planned comparisons were only conducted when
5
6 significant main effects or interactions (or trends, $p < 0.1$) were observed.
7
8

9 Results

10 Blood Glucose Levels

11
12 For mean blood glucose levels (\pm SE) see Figure 2. There was a main effect of
13
14 drink on blood glucose levels [$F(3,39)=59.42$, $p < 0.01$]. Post hoc analyses indicated
15
16 that the glucose drink lead to significantly higher blood glucose levels than the other
17
18 drinks (all p -values < 0.01). A main effect of time [$F(4,52)=37.43$, $p < 0.001$] was due
19
20 to increasing blood glucose levels over the course of the experimental session(all p -
21
22 values < 0.01) and a significant time*drink interaction [$F(12,156)=15.74$, $p < 0.001$]
23
24 showed that blood glucose levels were significantly higher at each post dose time
25
26 point following a glucose drink compared to any of the other treatments.
27
28
29
30
31
32

33 Cognitive Outcome Measures

34
35 Mean (\pm SD) baseline and post-drink change from baseline scores on the
36
37 cognitive outcome factors and VAS, on which significant drink effects were observed,
38
39 are displayed in Figures 3 and 4.
40
41
42

43 Power of Attention

44
45 The main effect of drink just failed to reach significance [$F(3,48)=2.30$,
46
47 $p=0.09$]. Planned comparisons between treatment drink and placebo demonstrated a
48
49 significant performance enhancement following glucose compared to placebo
50
51 ($t(48)=2.34$, $p < 0.05$) particularly 15-minutes post ingestion ($t(48)=-3.71$, $p < 0.01$). See
52
53 Figure 4(b).
54
55
56
57

58 Continuity of Attention

59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

There was a significant main effect of time [$F(3,48)=7.37, p<0.05$] whereby regardless of drink, performance was significantly worse 60-minutes post-drink than 15-minutes after drink ingestion ($p<0.05$).

Quality of Working Memory

A significant time*drink interaction was observed [$F(3,48)=3.34, p<0.05$]. Post hoc analyses revealed performance improvements 60-minutes after a fat drink compared to 15-minutes after a fat drink ($p<0.01$). Planned comparisons revealed that 15-minutes post drink, protein was associated with enhanced performance compared to placebo ($t(48)=2.15, p<0.05$) and 60-minutes post drink glucose was associated with impaired working memory compared to placebo ($t(48)=2.45, p<0.05$). See Figure 3(b).

Quality of Episodic, Secondary Memory

There was a significant main effect of drink [$F(3,48)=3.90, p<0.05$] with significantly better performance following a protein drink compared to a glucose drink ($p<0.05$). In addition, there was a significant time*drink interaction [$F(2.01, 32.18)=8.81, p<0.01$] which was due to the fact that performance was significantly better 60-minutes after a protein drink than after a fat or a glucose drink (both p -values <0.01). In addition, performance was enhanced 60-minutes following protein ingestion compared to placebo ingestion ($t(48)=4.42, p<0.001$). Furthermore, following a protein drink, performance improved significantly over time ($p<0.01$) whereas performance following a glucose drink deteriorated over time ($p<0.05$). See Figure 3(a).

Quality of Memory

There was a significant main effect of drink [$F(3,48)=4.969, p<0.01$]. Post hoc comparisons showed that performance was significantly better following a protein

1 drink than a glucose drink ($p < 0.05$). There was also a significant time*drink
2 interaction [$F(3,48)=8.14$, $p < 0.01$] with significantly enhanced performance 60-
3 minutes after a protein drink compared to performance following a fat drink ($p < 0.01$)
4 or a glucose drink ($p < 0.01$). Planned comparisons revealed that protein ingestion
5 significantly enhanced memory compared to placebo ($t(48)=2.49$, $p < 0.05$),
6 particularly at the 60-minutes post time point ($t(48)=4.72$, $p < 0.001$). In contrast, at 60-
7 minutes post ingestion glucose led to significantly impaired memory compared to
8 placebo ($t(48)=2.52$, $p < 0.05$). See Figure 3(c).

19 Speed of Memory

20 The time*drink interaction just missed significance [$F(3,48)=2.54$, $p=0.07$]. This
21 may be a result of one of the following observations: the slowest memory processing
22 was observed following the placebo drink, the glucose drink produced consistently
23 fast performance and speed following protein ingestion improved over time. Planned
24 comparisons revealed faster responses 15-minutes following glucose ingestion
25 ($t(48)=3.27$, $p < 0.05$) and following fat-ingestion ($t(48)=2.62$, $p < 0.05$) compared to
26 placebo. See Figure 3(d).

39 Combined Speed

40 The interaction of time*drink almost reached significance [$F(3,48)=2.70$,
41 $p=0.06$]. Planned comparisons revealed that 15-minutes post drink, fat and glucose
42 were associated with faster responses compared to placebo ($t(48)=2.77$, $p < 0.01$ and
43 $t(48)=3.85$, $p < 0.001$ respectively). See Figure 4(a).

52 Subjective Mood Measures

53 There was a main effect of time on alertness [$F(1,16)=7.23$, $p < 0.01$] with
54 participants reporting significantly higher alertness levels 10-minutes post drink
55
56
57
58
59
60
61
62
63
64
65

1 compared to either 15-minutes or 60-minutes post ingestion ($p < 0.05$ and $p < 0.01$,
2 respectively). See Figure 4(c).
3

4 <Figures 3 and 4 here>
5
6

7 Discussion

8
9 The present study further demonstrated the ability of macronutrients to affect
10 cognition and mood and that the effects are time-dependant and vary between
11 macronutrients. Protein facilitated working memory performance 15-minutes post
12 ingestion, and enhanced episodic memory 60-minutes post ingestion. Glucose
13 ingestion enhanced attentional processes (power of attention), speed of processing
14 (combined speed) and speed of memory 15-minutes post ingestion. Fat ingestion was
15 associated with enhanced speed of processing 15-minutes post drink. However,
16 glucose was also associated with impaired working memory (quality of working
17 memory) 60-minutes post ingestion.
18
19
20
21
22
23
24
25
26
27
28
29
30

31 Beneficial effects of protein (compared to placebo) were observed 15 and 60-
32 minutes post drink and were specifically targeting memory processes with enhanced
33 working memory 15-minutes post drink and enhanced episodic memory 60-minutes
34 post drink. Given the significant effects of protein on these memory factors, it is not
35 surprising that protein ingestion was also associated with enhanced memory accuracy
36 (quality of memory) as this factor is a combination of accuracy scores from all of the
37 memory tasks.
38
39
40
41
42
43
44
45
46
47

48 Beneficial effects following fat and glucose ingestion were also observed.
49 Fifteen minutes after fat ingestion cognitive processing speed (combined speed) and
50 speed of memory were faster than following placebo. Cognitive processing speed,
51 speed of memory and the ability to allocate attentional processes ('power of attention'
52 factor) were also faster 15-minutes following a glucose drink compared to placebo.
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Furthermore, 60-minutes after glucose ingestion working memory was significantly
2 impaired compared to placebo. This latter finding is not surprising since glucose is
3 metabolised quickly and enhancement is observed up to 20-minutes post ingestion
4 (e.g. Foster et al., 1998; Owens & Benton, 1994; Sünram-Lea et al., 2001; 2002a;
5 2002b; 2004) so enhancement at a later time point would not be expected. And indeed
6 impairments might be explained by a drop in blood glucose levels subsequent to
7 ingestion of a glucose load due to increased insulin output.
8
9

10
11
12
13
14
15
16
17 In general, the findings show beneficial effects on cognition soon after glucose
18 and fat ingestion (15-minutes post ingestion), whereas protein enhanced cognition at
19 later time points. There are some notable exceptions e.g. protein also enhanced
20 working memory 15-minutes post drink. These findings suggest that i) different
21 macronutrients have different windows of opportunity for performance improvements
22 and ii) different macronutrients target different cognitive domains. More specifically
23 it appears that protein has a beneficial effect on general memory processes, whereas
24 fat and glucose apparently target attentional processes and speed of processing.
25
26
27
28
29
30
31
32
33
34
35

36 In terms of glycaemic response the observed trajectories were nutrient-
37 dependent. As expected glucose ingestion led to significantly increased blood glucose
38 levels compared to fat, protein or placebo ingestion. This further supports the notion
39 that ingestion of energy, regardless of source, appears to improve certain aspects of
40 cognition and that at least some of the effects are independent of increases in blood
41 glucose levels (Kaplan et al., 2001).
42
43
44
45
46
47
48
49
50

51 The current findings are consistent with those of Kaplan et al. (2001) who
52 demonstrated macronutrient-mediated cognitive enhancement 15-minutes following
53 protein, glucose and fat. However, in contrast to the current findings, they also
54 observed maintained performance improvements following glucose ingestion 60-
55
56
57
58
59
60
61
62
63
64
65

1 minutes post ingestion. Moreover, the current study employed a repeated measures
2 design in order to reduce inter-participant variability. Steps were taken to minimise
3 the effects of drink order. Despite this a drink*time*drink order interaction was
4 observed on working memory. There were no main effects of drink order on any of
5 the measures. Future work should consider potential order effects perhaps studies
6 could employ between-participant designs.
7
8
9
10
11
12
13

14 The glucose facilitation effect of cognition has been widely reported in the
15 literature and has previously been reliably demonstrated with dosages of 25g glucose
16 in healthy, young adults (e.g. Foster et al., 1998; Kennedy & Scholey, 2000; Sünram-
17 Lea et al., 2001; 2002a; 2002b, 2004) and dosages of 25g and 50g in elderly adults
18 (e.g. Craft et al., 1994; Messier et al., 1997; Riby et al., 2004). The most robust
19 facilitation appears to be on memory (see reviews by Riby, 2004; Hoyland et al.,
20 2008); however glucose-mediated enhancement of other tasks including attention and
21 speed of processing (those that were enhanced in the current study) has also been
22 reported in previous research (e.g. reaction times: Owens & Benton, 1994;
23 information processing: Benton, Owens & Parker, 1994; Donohoe & Benton, 1999).
24 Taken together these findings suggest that glucose administration may not specifically
25 target memory processes and the effects of glucose may be more widespread.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Some previous papers have reported fat-mediated cognitive impairments (e.g.
44 Cunliffe, Obeid & Powell-Tuck, 1997; Kaplan et al., 2001; and Wells & Read, 1995).
45 However, Fischer et al. (2001) observed a beneficial effect of fat on a variety of
46 cognitive tasks which was not replicated in the current study. This could be due to the
47 optimal time-frame of fat metabolism which may have been missed as a result of the
48 relatively short experimental sessions employed in the current study (final testing was
49 60-minutes post ingestion). However, Fischer et al. (2001) observed faster reaction
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 times, improved short-term memory and improved attention at all time points (60-,
2 120- and 180 minutes) following fat ingestion. In the current study we observed faster
3 processing 15-minutes following a fat drink. The discrepancies between findings may
4 be a result of different dosages, time-frames and tasks that were employed by
5 different research groups. Further research is required to establish the effects of
6
7 different fat dosages on a variety of cognitive domains.
8
9
10
11
12
13

14 The current experiment attempted to separate nutritionally-mediated effects on
15 cognition and the effects of food perception on mood by assessing mood prior to
16 nutrient metabolism (10-minutes post drink) and 35-minutes and 80-minutes post-
17 drink. However, it is important to note that although the 10-minutes post drink time
18 point would precede absorption of both fat and protein, glucose would be
19 metabolically active so any observed effects could be nutritionally-mediated. Macht,
20 Gerer, & Ellgring (2003) found that increasing energy content of food was associated
21 with increased negative emotions and increased negative perceptions of the food (e.g.
22 more unhealthy and dangerous etc). However, in the current study no significant main
23 effects of drink type on self-rated alertness, contentment or calmness were observed
24 suggesting that energy content does not influence mood in healthy adults. However,
25 increased alertness was observed immediately post-drink, regardless of drink-type.
26 The fact that all drinks increased alertness suggests a general mechanism, possibly the
27 impact of hydration from all drinks. For example, Neave, Scholey & Emmett et al.
28 (2001) and Rogers, Kainth & Smit (2001) have shown that, compared to no drink,
29 water ingestion increases subjective alertness in both fasted and non-fasted, healthy,
30 young adults.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 Alternatively, the experimental situation may have been responsible for the
56 immediate post-drink increase in alertness observed. The 10-minutes post-drink mood
57
58
59
60
61
62
63
64
65

1 scale was administered in isolation whereas the other mood scales were all
2 administered immediately after the cognitive test battery. It may be the case that
3 participants felt significantly less alert at these times because they had just been
4 engaged, for 20-minutes, in cognitive tasks.
5
6
7
8

9 Macronutrient-specific mood effects demonstrated in previous research (e.g.
10 Fischer et al., 2001; Gibson et al., 1999) have not been replicated in the current study.
11 Fischer et al. (2001) found that carbohydrate ingestion led to an overall reduction in
12 depression scores on the POMS, which was not related to time of testing. Gibson et
13 al. found improved positive affect 120-minutes after a meal high in protein compared
14 to a meal low in protein. The observation of protein-mediated mood effects may
15 depend on a longer post-dose time-frame than that employed in the current study.
16 Alternatively, it may be that the mood scales used in the current study lacked
17 sensitivity to these effects, for example, previous research appears to demonstrate
18 specific improvement of subjective depression ratings associated with carbohydrate
19 ingestion (e.g. Fischer et al., 2001; Sayegh, Schiff, Wurtman, Spiers, McDermott &
20 Wurtman, 1995; Wurtman, Brzezinski, Wurtman, & Laferrere, 1989) but depression
21 was not measured in the current study. Moreover, Gibson administered a more
22 general mood scale and measured negative and positive affect. This was shown to be
23 sensitive to protein ingestion. Furthermore, Gibson did not administer a cognitive test
24 battery which in itself might affect mood.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 In terms of mechanisms, we can only speculate as biomarkers were not
49 measured in this study. There are, however, a number of potential mechanisms that
50 could be responsible for the effects of macronutrient ingestion on cognitive
51 performance and mood. Kaplan et al. (2001) suggest that carbohydrates, fat and
52 protein may improve performance via distinct mechanisms that are mediated by
53
54
55
56
57
58
59
60
61
62
63
64
65

1 different brain regions. Moreover, they argue that facilitation of certain aspects of
2 cognitive performance after administration of macronutrients which do not
3 significantly raise blood glucose levels suggest that facilitation of cognitive
4 performance might be due to more generic aspects of energy supply to the brain
5 (Kaplan et al., 2001). There is substantial evidence suggesting possible hippocampal
6 mediation for the glucose facilitation effect in both cognitive and physiological terms.
7 However, the fact that macronutrients which do not raise blood glucose levels also
8 improve certain aspects of cognitive functioning suggests that the enhancement effect
9 of certain foodstuffs on cognitive function may be nutrient-specific whereby the
10 action of glucose is on specific central mechanisms and other macronutrients have
11 their effects on more generalised peripheral mechanisms.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 It has been suggested that food-related memory enhancement may occur
28 through release of gastrointestinal peptides in response to ingestion of fat, protein,
29 and glucose (Flood et al., 1987; Flood & Morley, 1989). This notion is supported by
30 animal studies which have demonstrated that gastrointestinal peptides, such as
31 cholecystokinin (CCK; Flood & Morley, 1989; Flood et al., 1987), gastrin-releasing
32 peptide (Morley et al., 1994), and amylin (Flood & Morley, 1992) enhance memory
33 performance through vagus nerve stimulation. Moreover, enterostatin, an intestinal
34 peptide produced after food ingestion, has been shown to attenuate scopolamine-
35 induced amnesia (Takenaka, Nakamura, Jinsmaa, Lipowski, & Yoshikawa, 2001).
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Furthermore, Flood et al. (1987) and Morley et al. (1994) demonstrated that
50 CCK memory enhancement is abolished by vagotomy, suggesting that the central
51 effects of this peptide are mediated by activation of the ascending pathways of the
52 vagus nerve. In terms of mood effects, Cunliffe et al (1997) argue that the previously
53 observed fat mediated reduction in alertness and flicker fusion frequency could be
54
55
56
57
58
59
60
61
62
63
64
65

1 due to unspecified hormonal changes. Wells & Read (1996) attributed impaired
2 Bakkan performance to concomitant changes in mood and reduced alertness
3 following fat ingestion to increased CCK release stimulated by the presence of lipid
4 in the duodenum. CCK has been shown to increase sleepiness (e.g. Kapas et al., 1991;
5 Stacher, Bauer, & Steinringer, 1979). Thus CCK is attributed to both performance
6 enhancements and impairments. However, the findings reported in the current study
7 did not reveal fat-mediated performance impairments or mood effects. Alternatively,
8 cognitive enhancement could be related to circulating insulin. Previous research has
9 shown that insulin administration can enhance cognitive performance (e.g. Benedict
10 et al., 2004; Craft et al., 1999; Moosavi, Naghdi, Maghsoudi, & Zahedi, 2007). The
11 ingestion of protein and glucose is associated with increased levels of circulating
12 insulin (e.g. Nuttall, Mooradian, Gannon, Billington, & Krezowski, 1984) that could
13 potentially act directly on the CNS. However, this would not explain why fat-
14 mediated enhancements have been previously observed (Fischer et al., 2001) as fat
15 does not stimulate insulin secretion. Moreover, if insulin were responsible for
16 glucose- and protein-mediated cognitive enhancement the same or at least similar
17 effects would be observed following these two macronutrients.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41 The nutrient-specific profiles of peptide and hormone release are yet to be
42 elucidated and the effects of different peptides and peptide combinations on the CNS
43 and behaviour are not fully understood. However, it is possible that macronutrient-
44 specific peptide profiles may target different brain regions or neurotransmitter
45 systems.

46
47
48
49
50
51
52
53
54 In conclusion the current experiment has provided additional support for
55 nutritionally-mediated cognitive enhancement following the ingestion of
56 macronutrients, particularly 15-minutes post ingestion with sustained memory
57
58
59
60
61
62
63
64
65

1 enhancement 60-minutes after protein ingestion. Furthermore, the current findings
2 have revealed different temporal patterns of effects which suggest that the action of
3 different macronutrients on cognition may be related to nutrient-specific mechanisms.
4
5 However, a number of crucial questions remain to be answered before the beneficial
6 effects of macronutrient administration can be fully understood. For example, this
7 study has failed to clarify the basis of post-prandial mood effects and further
8 investigation is required. Moreover, there are other limitations which future studies
9 should aim to address. For example, we only investigated one dosage per
10 macronutrient and further research needs to be carried out to elucidate the effects of
11 different dosages of macronutrients on different cognitive domains. Moreover,
12 controlling for the potential confounding effects of cognitive testing on subjective
13 mood measures might help to elucidate macronutrient effects on mood. Finally, future
14 research should employ a longer post-dose period in order to further clarify the
15 window of opportunity of effects. In Western countries, the high incidence of obesity,
16 Type 2 diabetes and AD are associated with diet and increased fat intake (Martins et
17 al., 2006). The data suggests that modifiable lifestyle factors including diet may
18 contribute significantly to the risk of cognitive decline, including dementia.
19
20 Understanding the way nutrients affect behaviour will provide scientific evidence for
21 nutritional interventions aimed to increase health including optimal cognition and
22 psychological 'wellbeing'.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 References

52
53 Benedict, C., Hallschmid, M., Hatke, B., Fehm, H.L., Born, J. & Kern, W. (2004).
54
55 Intranasal insulin improves memory in humans. *Psychoneuroendocrinology*,
56
57 29(10), 1326-1334.
58
59
60
61
62
63
64
65

- 1 Benton, D., Owens, D.S. & Parker, P.Y. (1994). Blood glucose influences memory
2 and attention in young adults. *Neuropsychologia*, 32(5), 595-607.
3
- 4 Benton, D., Ruffin, M-P., Lassel, T., Nabb, S., Messaoudi, M. & Vinoy, S., Desor, D.,
5 & Lang, V. (2003) The delivery rate of dietary carbohydrates affects cognitive
6 performance in both rats and humans. *Psychopharmacology*, 166, 86–90.
7
- 8 Benton, D. & Sargent., J. (1992). Breakfast, blood glucose and memory. *Biological*
9 *Psychology*, 33(2-3), 207-210.
10
- 11 Bond, A. & Lader, M. (1974). The use of analogue scales in rating subjective feelings.
12 *British Journal of Medical Psychology*, 47(3), 211-218.
13
- 14 Craft, S., Murphy, C., & Wemstrom, J. (1994). Glucose effects on complex memory
15 and nonmemory tasks: the influence of age, sex and glucoregulatory response.
16 *Psychobiology*, 22, 95-105.
17
- 18 Craft, S., Asthana, S., Newcomer, J.W., Wilkinson, C.W., Matos, I.T., & Baker, L.D.
19 et al. (1999). Enhancement of memory in Alzheimer’s disease with insulin and
20 Somatostatin, but not glucose. *Archives of General Psychiatry*, 56, 1135-1140.
21
- 22 Cunliffe, A., Obeid, O.A., & Powell-Tuck, J. (1997). Post-prandial changes in
23 measures of fatigue: effect of a mixed or a pure carbohydrate or pure fat meal.
24 *European Journal of Clinical Nutrition*, 51, 831-838.
25
- 26 Donohoe, R.T. & Benton, D. (1999). Cognitive functioning is susceptible to the level
27 of blood glucose. *Psychopharmacology*, 145, 378-385.
28
- 29 Erdfelder, E., Faul, F., & Buchner, A. (1996). GPOWER: A general power analysis
30 program. *Behavior Research Methods, Instruments, & Computers*, 28, 1-11.
31
- 32 Fischer, K., Colombani, P.C., Langhans, W., & Wenk, C. (2001). Cognitive
33 performance and its relationship with postprandial metabolic changes after
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 ingestion of different macronutrients in the morning. *British Journal of*
2 *Nutrition*, 85, 393-405.
3

4 Flood, J.F. & Morley, J.E. (1989). Cholecystokinin receptors mediate enhanced
5 memory by feeding and gastrointestinal peptides. *Peptides*, 10, 808-813.
6

7 Flood, J.F., & Morley, J.E. (1992). Differential effects of amylin on memory
8 processing using peripheral and central routes of administration. *Peptides*,
9 13(3), 577-580.
10

11 Flood, J.F., Smith, G.E., & Morley, J.E. (1987). Modulation of memory processing
12 by Cholecystokinin: dependence on the vagus nerve. *Science*, 236, 832-834.
13

14 Foster, J.K. Lidder, P.G., & Sünram, S.I. (1998). Glucose and memory: fractionation
15 of enhancement effects? *Psychopharmacology*, 137, 259-270.
16

17 Gibson, E. L., Checkley, S., Popadopoulos, A., Poon, L., Daley, S., & Wardle, J.
18 (1999). Increased salivary cortisol reliably induced by a protein-rich midday
19 meal. *Psychosomatic Medicine*, 61, 214-224.
20

21 Gilsenan, M.B., de Bruin, E.A., & Dye, L. (2009). The influence of carbohydrate on
22 cognitive performance: a critical evaluation from the perspective of glycaemic
23 load. *British Journal of Nutrition*, 100, 941-949.
24

25 Hoyland, A., Lawton, C.L., & Dye, L. (2008). Acute effects of macronutrient
26 manipulations on cognitive test performance in healthy young adults: A
27 systematic research review. *Neuroscience and Biobehavioural Reviews*, 32(1),
28 72-85.
29

30 Ingwersen, J., Defeyter, M.A., Kennedy, D.O., Wesnes, K.A., & Scholey, A.B.
31 (2007). A low glycaemic index breakfast cereal preferentially prevents
32 children's cognitive performance from declining throughout the morning.
33 *Appetite*, 49(1), 240-244.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Jones, E., Sünram-Lea S.I. & Threadgold, E. (2005). The effects of protein and
2 glucose ingestion on cognitive performance and mood. *Appetite*, 45(3), 362-
3 377. In Abstracts of the joint meeting of the British Feeding and Drinking
4 Group (BFDG), and the Werkgroep Voedingsgewoonten (WEVO; Dutch
5 Working Group on Food Habits) 11–12 April 2005 Wageningen University,
6 The Netherlands.

7
8
9
10
11
12
13
14 Kapas, L., Obal, F., Ferkas, I., Payne, L.C., Sary, G., Rubicsek, G., Krueger, J.M.
15 (1991). Cholecystokinin promotes sleep and reduces food intake in diabetic
16 rats. *Physiology of Behaviour*, 50, 417-420.

17
18
19
20
21
22 Kaplan, R.J., Greenwood, C.E., Winocur, G., & Wolever, T.M.S. (2000). Cognitive
23 performance is associated with glucose regulation in healthy, elderly persons
24 and can be enhanced with glucose and dietary carbohydrates. *American*
25 *Journal of Clinical Nutrition*, 72, 825-836.

26
27
28
29
30
31 Kaplan, R.J., Greenwood, C.E., Winocur, G., & Wolever, T.M.S. (2001). Dietary
32 protein, carbohydrate and fat enhance memory performance in the healthy
33 elderly. *American Journal of Clinical Nutrition*, 74, 687-693.

34
35
36
37
38
39 Kennedy, D.O., & Scholey, A.B. (2000). Glucose administration, heart rate and
40 cognitive performance effects of increasing mental effort.
41 *Psychopharmacology*, 149, 63-71.

42
43
44
45
46 Kennedy, D.O., Scholey, A.B., Tildsley, N.T.J., Perry, E.K. and Wesnes, K.A. (2002).
47 Modulation of mood and cognitive performance following acute
48 administration of *Melissa officinalis* (lemon balm). *Pharmacology,*
49 *Biochemistry and Behaviour*, 72, 953-964.

50
51
52
53
54
55
56 Kennedy, D.O., Scholey, A.B. and Wesnes, K.A. (2001) Dose dependent
57
58
59
60
61
62
63
64
65

1 changes in cognitive performance and mood following acute administration of
2 Ginseng to healthy young volunteers. *Nutritional Neuroscience*, 4, 295-310.

3
4 Kennedy, D.O., Scholey, A.B. and Wesnes, K.A. (2002) Modulation of
5
6 cognition and mood following administration of single doses of Ginkgo
7
8 biloba, Ginseng and a Ginkgo/Ginseng combination to healthy young adults.
9
10
11
12 *Physiology and Behaviour* 75,1-13.

13
14 Macht, M., Gerer, J., & Ellgring, H. (2003). Emotions in overweight and normal-
15
16 weight women immediately after eating foods differing in energy. *Physiology*
17
18 and *Behaviour*, 80, 367-374.

19
20
21 Mahoney, C.R., Taylor, H.A., Kanarek, R.B. & Samuel, P. (2005). Effect of breakfast
22
23 composition on cognitive processes in elementary school children. *Physiology*
24
25 and *Behaviour*, 85, 635–645.

26
27
28
29 Martins, I.J, Hone, E., Foster, J.K., Sünram-Lea, S.I., Gnjec, A., Martins, R.N.,
30
31 Gandy, S. (2006). Cholesterol and Diabetes as risk factors for Alzheimer's
32
33 Disease. *Molecular Psychiatry*, 11, 721-736.

34
35
36 Messier, C. (2004). Glucose improvement of memory: a review. *European Journal of*
37
38 *Pharmacology*, 490 (1-3), 33-57.

39
40
41 Messier, C., Gagnon, M. & Knott, V. (1997). Effect of glucose and peripheral glucose
42
43 regulation on memory in the elderly. *Neurobiology of Aging*, 18 (3), 297-304.

44
45
46 Moosavi, M., Naghdi, N., Maghsoudi, N., & Zahedi Asl, S. (2007). Insulin protects
47
48 against stress-induced impairments in water maze performance. *Behavioural*
49
50 *Brain Research*, 176, 230-236.

51
52
53
54 Morley J.E., Flood, J.F., Silver, A.J., Kaiser, F.E. (1994). Effects of peripherally
55
56 secreted hormones on behaviour. *Neurobiology of Aging*, 15(4), 573-577

57
58
59 Neave N, Scholey AB, Emmett JR, Moss M, Kennedy DO & Wesnes KA (2001)

60
61
62
63
64
65

Water ingestion improves subjective alertness, but has no effect on cognitive performance in dehydrated healthy young volunteers. *Appetite* 37:255-256.

Nuttall, F.Q., Mooradian, A.D., Gannon, M.C., Billington, C., & Krezowski, P. (1984). Effect of protein ingestion on the glucose and insulin response to a standardized oral glucose load. *Diabetes Care*, 7(5), 465-470.

Owens, D.S., & Benton, D. (1994). The impact of raising blood glucose levels on reaction time. *Neuropsychobiology*, 30, 106-113.

Riby, L.M. (2004). The impact of age and task domain on cognitive performance: a meta-analytic review of the glucose facilitation effect. *Brain Impairment*, 5(2), 145-165.

Riby, L.M., Laws, A.S., McLaughlin, J., & Murray, J. (2011). Preliminary evidence that glucose ingestion facilitates prospective memory performance. *Nutrition Research*, 31(5), 370-377.

Riby, L.M., Meikle, A., & Glover, C. (2004). The effects of age, glucose ingestion and gluco-regulatory control on episodic memory. *Age and Ageing*, 33, 483-487.

Rogers, P.J., Kainth, A., & Smit, H.J. (2001). A drink of water can improve or impair mental performance depending on small differences in thirst. *Appetite*, 36, 57-58.

Scholey, A.B., Harper, S., & Kennedy, D.O. (2001). Cognitive demand and blood glucose. *Physiology and Behaviour*, 73(4), 585-592.

Scholey, A.B., Sünram-Lea, S.I., Greer, J., Elliott, J., & Kennedy, D.O. (2009). Glucose administration prior to a divided attention task improves tracking performance but not word recognition: evidence against differential memory enhancement? *Psychopharmacology*, 202, 549-558.

- 1 Sayegh, R., Schiff, I., Wurtman, J., Spiers, P., McDermott, J. & Wurtman, R. (1995).
2 The effect of a carbohydrate-rich beverage on mood, appetite, and cognitive in
3 women with premenstrual syndrome. *Obstetrics & Gynaecology*, 86(4), 520-
4 528.
5
6
7
8
9
10 Smith, M.A. & Foster, J.K. (2008). Glucoregulatory and order effects on verbal
11 episodic memory in healthy adolescents after oral glucose administration.
12 *Biological Psychology*, 79, 209-215.
13
14
15
16
17 Smith, M.A., Hii, H.L., Foster, J.K., & van Eekelen, J.A.M. (2009). Glucose
18 enhancement of memory is modulated by trait anxiety in healthy adolescent
19 males. *Journal of Psychopharmacology*; 0: 0269881109348164v1
20
21
22
23
24 Stacher, G., Bauer, H., & Steinringer, H. (1979). Cholecystokinin decreases appetite
25 and activation evoked by stimuli arising from the preparation of a meal in
26 man. *Physiology of Behaviour*, 23, 325-331.
27
28
29
30
31
32 Sünram-Lea, S.I., Birchall, R.J., Wesnes, K.A., & Petrini, O. (2004). The effect of
33 acute administration of 400mg of Panax Ginseng on cognitive performance
34 and mood in healthy young volunteers. *Current Topics in Nutraceutical*
35 *Research*, 3(1), 251-254.
36
37
38
39
40
41 Sünram-Lea, S.I., Foster, J.K., Durlach, P., & Perez, C. (2001). Glucose facilitation of
42 cognitive performance in healthy young adults: examination of the influence
43 of fast-duration, time of day and pre-consumption plasma glucose levels.
44 *Psychopharmacology*, 157, 46-54.
45
46
47
48
49
50
51 Sünram-Lea, S.I., Foster, J.K., Durlach, P., & Perez, C. (2002a). The effect of
52 retrograde and anterograde glucose administration on memory performance in
53 healthy young adults. *Behavioural Brain Research*, 134, 505-516.
54
55
56
57
58
59 Sünram-Lea, S.I., Foster, J.K., Durlach, P., & Perez, C. (2002b). Investigation into the
60
61
62
63
64
65

1 significance of task difficulty and divided allocation of resources on the
2 glucose memory facilitation effect. *Psychopharmacology*, 160, 387-397.
3

4 Sünram-Lea, S.I., Foster, J.K., Durlach, P., & Perez, C. (2004). The influence of fat
5 co-administration on the glucose memory facilitation effect. *Nutritional*
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Neuroscience, 1, 21-32.

Takenaka, Y., Nakamura, F., Jinsmaa, Y., Lipowski, A.W., & Yoshikawa, M. (2001).
Enterostatin (VPDPR) has anti-analgesic and anti-amnesic activities.
Bioscience, Biotechnology and Biochemistry, 65(1), 236-238.

Wells, A.S., Read, N.W., & Craig, A. (1995). Influences of dietary and intraduodenal
lipid on alertness, mood and sustained concentration. *British Journal of*
Nutrition, 74, 115-123.

Wells, A.S., & Read, N.W. (1996). Influences of fat, energy and time of day on mood
and performance. *Physiology and Behaviour*, 59(6), 1069-1076.

Wesnes, K.A., Faleni, R.A., Hefting, N.R., Hoogsteen, G., Houben, J.J.G.,
Jenkins, E., Jonkman, J.H.G., Leonard, J., Petrini, O. and van Lier, J.J.
(1997). The cognitive, subjective, and physical effects of a Ginkgo
biloba/Panax Ginseng combination in healthy volunteers with neurasthenic
complaints. *Psychopharmacology Bulletin* 33, 677-683.

Wesnes, K.A., Pincock, C., Richardson, D. Helm, G., & Hails, S. (2003). Breakfast
reduces declines in attention and memory over the morning in schoolchildren.
Appetite, 41, 329-331.

Wesnes, K.A., Ward, T., Ayre, G. and Pincock, C. (1999) Validity and utility of the
cognitive drug research (CDR) computerised assessment system: a review
following fifteen years of usage. *European Neuropsychopharmacology* 9,
S368.

1 Wesnes, K. A., Ward, T., McGinty, A., & Petrini, O. (2000). The memory enhancing
2 effects of a Gingko biloba/Panax ginseng combination in healthy middle-aged
3
4
5 volunteers. *Psychopharmacology*, 152, 353-361.
6

7 Wurtman, J.J., Brzezinski, A., Wurtman, R.J., & Laferrere, B. (1989). Effect of
8
9
10 nutrient intake on premenstrual depression. *American Journal of Obstetrics*
11
12 and *Gynaecology*, 161(5), 1228-1234.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Composition of treatment drinks

Glucose	Fat	Protein	Placebo
40g glucose dextrose powder (Thornton and Ross Ltd, Huddersfield, UK)	16g Pura Vegetable oil (supplied by Sainsbury's Plc, UK)	40g Casilan 90% protein powder (supplied by Boots Plc., UK) ^a	2g aspartame (Candarel, Merisant UK Ltd)
10ml Lemon juice	10ml Lemon juice	10ml Lemon juice	10ml Lemon juice
260 ml water	249 ml water	260ml water	290ml water

^a The other 10% is made up of small amounts of carbohydrate, fat, fibre, sodium and calcium.

Table 2. Mean (\pm SD) Cognitive Performance Scores at Baseline and the Post-drink

Change from Baseline


Measure	Baseline score	Post-drink change from baseline	
		15-minutes	60-minutes
Immediate word recall (% accuracy)			
Placebo	35.93 (11.35)	0.00 (13.54)	-4.12 (16.65)
Fat	37.78 (14.99)	-1.57 (13.29)	-6.67 (13.69)
Protein	42.78 (13.00)	-5.10 (15.42)	-4.31 (13.63)
Glucose	43.14 (10.24)	-7.26 (10.82)*	-11.57 (12.91)*
Total	39.91 (12.40)	-3.48 (13.27)	-6.67 (5.90)
Delayed word recall (% accuracy)			
Placebo	24.63 (11.15)	-10.39 (13.64)	-7.65 (16.06)
Fat	25.74 (14.09)	-10.78 (10.37)	-16.08 (14.96)
Protein	28.70 (16.01)	-13.53 (14.12)	-14.31 (14.56)
Glucose	25.29 (9.79)	-10.98 (10.72)	-14.12 (8.86)
Total	26.09 (12.76)	-11.42 (12.13)	-13.04 (13.61)
Simple reaction time (ms)			
Placebo	265.99 (23.92)	16.54 (25.31)	18.35 (23.86)
Fat	266.89 (33.99)	11.29 (25.05)	16.05 (22.90)
Protein	267.71 (39.18)	16.23 (23.72)	17.78 (18.97)
Glucose	265.30 (28.74)	5.56 (18.73)	10.78 (22.62)
Total	266.47 (31.46)	12.41 (23.20)	15.74 (22.09)
Choice reaction time accuracy (%)			
Placebo	94.56 (4.94)	0.00 (3.46)	-1.41 (3.80)
Fat	95.11 (4.66)	-2.24 (6.08)	-0.71 (4.36)
Protein	94.78 (5.49)	0.59 (3.52)	-0.12 (2.78)
Glucose	93.41 (6.96)	0.59 (4.35)	2.00 (4.53)
Total	94.47 (5.51)	-0.265 (4.35)	-0.06 (3.87)
Choice reaction time (ms)			
Placebo	398.29 (41.25)	4.56 (36.37)	-2.94 (33.32)
Fat	397.22 (49.16)	-5.46 (23.63)	-1.83 (30.19)
Protein	391.08 (42.22)	1.15 (26.69)	12.80 (31.26)
Glucose	391.81 (45.94)	-7.70 (25.63)	9.11 (27.39)
Total	394.60 (44.64)	-1.86 (28.08)	4.29 (30.54)
Digit vigilance accuracy (%)			
Placebo	94.82 (8.14)	0.91 (6.76)	-2.48 (5.26)
Fat	95.19 (5.62)	0.39 (4.98)	-3.66 (5.33)
Protein	95.31 (4.81)	0.52 (4.75)	-0.52 (4.87)
Glucose	95.95 (4.66)	-0.65 (5.32)	-2.75 (6.16)
Total	95.32 (5.81)	0.29 (5.45)	-2.35 (5.41)
Digit vigilance reaction time (ms)			
Placebo	423.18 (47.70)	17.43 (38.98)	11.23 (26.17)
Fat	412.15 (36.04)	11.09 (27.93)	15.07 (31.08)
Protein	417.99 (38.75)	14.86 (33.57)	13.37 (29.16)
Glucose	433.48 (41.02)	-8.81 (26.08)	-7.31 (34.13)
Total	421.70 (40.88)	8.643 (31.64)	8.09 (30.14)
Spatial memory original item accuracy (%)			
Placebo	93.40 (6.60)	-2.57 (13.08)	1.10 (10.65)
Fat	94.10 (6.24)	-2.21 (11.04)	-0.37 (7.48)
Protein	88.89 (13.14)	6.25 (15.93)	4.78 (12.01)
Glucose	94.12 (5.62)	-1.84 (7.25)	-5.15 (15.51)
Total	92.63 (7.90)	-0.09 (11.83)	0.09 (11.41)
Spatial memory new item accuracy (%)			
Placebo	97.22 (4.28)	-1.76 (8.09)	-3.24 (9.99)
Fat	98.89 (2.14)	-3.82 (5.74)	-2.35 (4.72)
Protein	92.78 (11.14)	2.06 (12.00)	3.53 (13.32)
Glucose	95.88 (5.93)	2.35 (4.37)	-3.82 (10.83)
Total	96.19 (5.87)	-0.29 (7.55)	-1.47 (9.72)
Spatial memory sensitivity index (SI)			
Placebo	0.91 (0.09)	-0.04 (0.19)	-0.02 (0.19)
Fat	0.94 (0.06)	-0.06 (0.13)	-0.03 (0.05)
Protein	0.82 (0.23)	0.08 (0.28)	0.08 (0.24)
Glucose	0.90 (0.88)	0.01 (0.07)	-0.09 (0.23)
Total	0.89 (0.32)	-0.0025 (0.17)	-0.015 (0.18)
Spatial memory reaction time (ms)			
Placebo	564.36 (127.08)	-22.62 (60.40)	-37.32 (53.37)
Fat	586.87 (151.34)	-62.16 (93.23)	-54.58 (107.43)
Protein	599.05 (157.44)	-41.68 (167.28)	-76.87 (152.22)
Glucose	616.11 (191.16)	-73.34 (153.69)	-86.22 (93.94)
Total	591.60 (156.76)	-49.95 (118.65)	-63.748 (101.74)

Measure	Baseline score	Post-drink change from baseline	
		15-minutes	60-minutes
Numeric working memory original item accuracy (%)			
Placebo	87.28 (13.55)	-1.18 (8.32)	-0.78 (7.20)
Fat	88.27 (14.09)	-2.22 (5.27)	0.13 (7.18)
Protein	87.78 (11.79)	-2.09 (6.00)	-1.31 (5.21)
Glucose	88.76 (14.18)	-3.40 (7.08)	-2.88 (6.61)
Total	88.02 (13.40)	-2.22 (6.67)	-1.21 (6.55)
Numeric working memory new item accuracy (%)			
Placebo	94.08 (9.88)	1.57 (5.14)	1.31 (4.58)
Fat	92.96 (9.50)	-0.13 (5.00)	2.61 (4.86)
Protein	94.57 (11.36)	0.00 (5.21)	-0.65 (6.61)
Glucose	93.60 (8.16)	1.18 (3.69)	-1.05 (4.85)*
Total	93.80 (9.73)	0.66 (4.76)	0.56 (5.23)
Numeric working memory sensitivity index (SI)			
Placebo	0.82 (0.22)	0.01 (0.10)	0.01 (0.09)
Fat	0.82 (0.23)	-0.02 (0.08)	0.03 (0.10)
Protein	0.83 (0.22)	-0.16 (0.09)	-0.02 (0.11)
Glucose	0.83 (0.21)	-0.02 (0.72)	-0.04 (0.08)
Total	0.83 (0.22)	-0.05 (0.25)	-0.005 (0.10)
Numeric working memory speed (ms)			
Placebo	546.67 (91.19)	-16.48 (79.04)	-29.14 (63.00)
Fat	530.86 (72.16)	-19.87 (43.64)	-22.71 (57.97)
Protein	532.73 (87.55)	-7.78 (37.66)	-37.78 (33.48)
Glucose	557.27 (92.22)	-37.31 (33.98)	-37.92 (72.86)
Total	541.88 (85.78)	-20.36 (48.58)	-31.89 (56.83)
Word recognition original item accuracy (%)			
Placebo	60.00 (16.01)	1.96 (12.19)	-0.39 (13.01)
Fat	55.55 (16.80)	4.31 (17.79)	1.18 (19.75)
Protein	61.48 (17.04)	-3.14 (15.83)	-1.57 (17.08)
Glucose	55.69 (18.25)	2.35 (11.29)	-4.71 (12.64)
Total	58.18 (17.03)	1.37 (14.28)	-1.37 (15.62)
Word recognition new item accuracy (%)			
Placebo	87.78 (10.30)	-1.57 (10.94)	-5.88 (10.24)
Fat	86.67 (14.64)	-0.79 (16.31)	-7.84 (15.14)
Protein	86.30 (13.23)	1.57 (8.67)	-6.67 (12.91)
Glucose	84.31 (16.32)	-2.75 (13.96)	-6.67 (11.55)
Total	86.27 (13.62)	-0.89 (12.47)	-6.77 (12.46)
Word recognition sensitivity index (SI)			
Placebo	0.53 (0.17)	0.00 (0.18)	-0.08 (0.19)
Fat	0.51 (0.21)	0.01 (0.30)	-0.11 (0.32)
Protein	0.53 (0.24)	0.01 (0.14)	-0.10 (0.21)
Glucose	0.49 (0.23)	-0.05 (0.21)	-0.16 (0.21)
Total	0.52 (0.21)	-0.008 (0.21)	-0.11 (0.23)
Word recognition reaction time (ms)			
Placebo	651.31 (129.45)	64.53 (241.18)	-21.01 (78.99)
Fat	616.47 (70.52)	-5.89 (74.50)	17.73 (88.08)
Protein	636.27 (101.26)	-2.63 (93.60)	-23.55 (85.32)
Glucose	622.28 (83.61)	13.96 (78.53)	19.30 (101.59)
Total	631.58 (96.21)	17.49 (121.95)	-1.88 (88.50)
Picture recognition original item accuracy (%)			
Placebo	75.00 (17.66)	0.29 (13.40)	1.76 (9.67)
Fat	77.22 (20.95)	1.18 (16.25)	-5.88 (16.23)
Protein	80.83 (17.00)	-4.41 (12.98)	-5.00 (17.05)
Glucose	80.59 (15.19)	-5.00 (15.91)	-5.29 (12.05)
Total	78.41 (17.70)	-1.99 (14.64)	-3.60 (13.75)
Picture recognition new item accuracy (%)			
Placebo	91.11 (6.08)	-3.24 (9.34)	-1.47 (8.62)
Fat	88.61 (11.09)	2.35 (12.13)	-0.29 (10.07)
Protein	90.00 (9.24)	0.29 (6.27)	-2.94 (6.14)
Glucose	89.41 (8.08)	-2.06 (12.63)	0.59 (9.50)
Total	89.78 (8.62)	-0.67 (10.09)	-1.03 (8.58)


Measure	Baseline score	Post-drink change from baseline	
		15-minutes	60-minutes
Picture recognition sensitivity index (SI)			
Placebo	0.69 (0.17)	-0.04 (0.17)	0.00 (0.13)
Fat	0.67 (0.29)	0.048 (0.21)	-0.05 (0.21)
Protein	0.72 (0.24)	-0.03 (0.14)	-0.08 (0.16)
Glucose	0.72 (0.17)	-0.07 (0.21)	-0.04 (0.13)
Total	0.70 (0.22)	-0.02 (0.18)	-0.04 (0.16)
Picture recognition reaction time (ms)			
Placebo	718.72 (91.13)	23.15 (60.94)	0.38 (48.44)
Fat	709.04 (87.31)	-1.38 (67.94)	-6.70 (74.57)
Protein	705.53 (105.12)	41.10 (153.71)	-1.82 (76.48)
Glucose	725.84 (87.80)	-26.72 (76.48)	-17.75 (85.09)
Total	714.78 (92.84)	9.04 (89.77)	-6.47 (71.15)

Difference between treatment and placebo significant at: * 0.05 level and ** 0.01 level


Cognitive Outcome Measure




Figure(s)




(a) Quality of Secondary Memory




(b) Quality of Working Memory




(c) Quality of Memory




(d) Speed of Memory



(a) Combined Speed



(b) Power of Attention



(c) Self-rated Alertness




Figure 1. Schematic representation of the CDR battery showing the cognitive tasks, individual task outcome measures and the factors derived by factor analysis. Arrows indicate that a task outcome measure contributes to the given factor: Power of Attention, Continuity of Attention, Quality of Memory and Speed of Memory. Format of figure taken from Kennedy, Scholey, Tildesley, Perry and Wesnes (2002).

Figure 2. Mean Blood Glucose Levels over the Course of the Experimental Sessions

Figure 3. Profile of effects of macronutrients on cognitive outcome measures relating to memory: a) Quality of secondary memory, b) Quality of working memory, c) Quality of memory and c) Speed of memory at the two post dose time points (15-minutes and 60-minutes). Planned comparisons revealed significant enhancement following protein ingestion, compared to placebo on Quality of secondary memory (60-minutes post, $p < 0.001$), Quality of working memory (15-minutes post, $p < 0.05$) and quality of memory (60-minutes post, $p < 0.001$). Glucose ingestion, compared to placebo: impaired working memory (60-minutes post drink, $p < 0.05$), impaired quality of memory (60-minutes post, $p < 0.05$) and led to faster memory processing (speed of memory, 15-minutes post, $p < 0.05$). Fat ingestion led to faster memory processing (speed of memory) compared to placebo (15-minutes post, $p < 0.05$).

Figure 4. Profile of cognitive factors: a) Combined Speed and b) Power of Attention and c) self rated mood (measured by VAS). Power of attention was significantly improved following glucose compared to placebo (particularly 15-minutes post, $p < 0.01$). Fat and glucose led to faster overall responses (combined speed) 15-minutes post drink, compared to placebo ($p < 0.01$ and $p < 0.001$ respectively). Self-rated alertness was significantly higher 10-minutes post drink, regardless of drink, than 15- and 60-minutes post drink ($p < 0.05$ and $p < 0.01$ respectively).