

Northumbria Research Link

Citation: Easthaugh, Susan, Bradley, Gemma, Pee, Lorna and Donnelly, Joanna (2019) Occupational therapy-led pulmonary rehabilitation : a practice analysis. *British Journal of Occupational Therapy*, 82 (12). pp. 770-774. ISSN 0308-0226

Published by: UNSPECIFIED

URL:

This version was downloaded from Northumbria Research Link: <http://northumbria-test.eprints-hosting.org/id/eprint/53075/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



UniversityLibrary



Northumbria
University
NEWCASTLE



Occupational Therapy-Led Pulmonary Rehabilitation: A Practice Analysis

Journal:	<i>British Journal of Occupational Therapy</i>
Manuscript ID	110-May-2019-PA.R1
Manuscript Type:	Practice Analysis
Key Areas:	Adult Physical < Clinical, Assessment < Clinical, Professional Development
Keywords:	Pulmonary Rehabilitation, Chronic Obstructive Pulmonary Disease, Occupational Therapy
Abstract:	<p>Statement of context: Pulmonary Rehabilitation (PR) is accepted as an essential strategy for the management of respiratory disease. However, there is wide variation in models of service delivery and evidence to understand which elements are most effective is less clear. This analysis outlines key elements of an occupational therapy-led PR programme.</p> <p>Critical reflection on practice: It is proposed that the core focus on occupation, alongside the unique skills and reasoning of occupational therapists, contribute to positive outcomes for service users.</p> <p>Implications for practice: Occupational therapists are ideally placed to lead and develop PR programmes. As there is genuine uncertainty about which elements are most effective, this analysis should encourage further evaluation and research.</p>

Occupational Therapy-Led Pulmonary Rehabilitation: A Practice Analysis

Abstract

Statement of context: Pulmonary Rehabilitation (PR) is accepted as an essential strategy for the management of respiratory disease. However, there is wide variation in models of service delivery and evidence to understand which elements are most effective is less clear. This analysis outlines key elements of an occupational therapy-led PR programme.

Critical reflection on practice: It is proposed that the core focus on occupation, alongside the unique skills and reasoning of occupational therapists, contribute to positive outcomes for service users.

Implications for practice: Occupational therapists are ideally placed to lead and develop PR programmes. As there is genuine uncertainty about which elements are most effective, this analysis should encourage further evaluation and research.

Key Words

Pulmonary Rehabilitation, Chronic Obstructive Pulmonary Disease, Occupational Therapy

Statement of context

Chronic Obstructive Pulmonary Disease (COPD) is a chronic lung condition which causes multiple symptoms affecting quality of life and is the third largest cause of mortality worldwide (Department of Health (DH), 2012). It is estimated that 1.2 million people are living with COPD in the UK, over 170 million people living with the condition worldwide, and there is general acceptance that these statistics will underestimate the prevalence of the condition due to undiagnosed and mis-diagnosed cases, and issues with definitions and classifications (British Lung Foundation (BLF), 2017; Adeloje et al, 2015).

COPD is an incurable condition with significant personal and economic costs. It is estimated that lung disease costs the UK over £11 billion each year in costs to the NHS, social care and in lost productivity (BLF, 2017). Whilst technological advances, alongside public health drivers, are influencing mortality and morbidity rates for other major groups of health conditions (for example cardiovascular conditions and cancers), COPD is the only major cause of death that is increasing (BLF, 2017). The prevalence amongst the population, alongside the burden and cost at individual and societal levels, means understanding ways to manage this condition is a clear priority.

Pulmonary Rehabilitation (PR) is defined as a "comprehensive intervention based on a thorough patient assessment followed by patient tailored therapies that include, but are not limited to, exercise training, education, and behaviour change, designed to improve the physical and psychological condition of people with chronic respiratory disease and to promote the long-term adherence to health-enhancing behaviours" (Spruit et al, 2013). There is well-established evidence

1
2
3 that PR improves health-related quality of life for people with COPD and it is therefore accepted as
4 an essential strategy for people living with this condition (Global Initiative for Chronic Obstructive
5 Lung Disease (GOLD), 2017; McCarthy et al, 2015). However, whilst general efficacy is accepted, it is
6 still unclear what components of PR lead to the best outcomes for patients, which includes
7 uncertainty over length, frequency, location and professional involvement (McCarthy et al, 2015).
8
9

10 11 12 **Critical Reflection on Practice**

13
14 This practice analysis reflects on the development of an occupational therapy-led model of PR within
15 the North East of England. The model was first trialled for one year as part of a pilot project with one
16 identified GP practice and following successful evaluation, was recommissioned on an ongoing basis
17 and is now delivered across three locality areas. Key features of the programme include: it is led, and
18 primarily facilitated, by occupational therapists; it is delivered mainly in community leisure facilities;
19 it adopts a cohort model where individuals start and finish together for the 6-week programme and;
20 it utilises existing local authority services and personnel to deliver the exercise component of the
21 programme.
22
23
24

25
26 Despite a lack of evidence to support specific models of service delivery, physiotherapy-led PR
27 services have emerged as more common (Yohannes & Connelly, 2004) and reviews of published
28 research yield very few studies which specifically mention occupational therapy (Walker et al, 2016).
29 One likely explanation is linked to the centrality of exercise and strengthening as a key element of PR
30 programmes and a central domain of concern for physiotherapists, yet there is limited evidence
31 which supports this as the central focus, or supports the optimum composition of teams. In order to
32 further understanding of the contribution of occupational therapy in this area, this analysis
33 highlights some of the key features of the occupational-therapy led programme, alongside a
34 reflective discussion of how these elements have been developed in line with the core values and
35 philosophy of occupational therapy.
36
37
38
39
40
41
42

43 *The Occupation-Focussed Initial Assessment*

44 **As part of this occupational therapy-led PR programme,** the initial assessment is conducted in the
45 person's own home and includes standardised and non-standardised elements. The British Thoracic
46 Society Quality Standards for Pulmonary Rehabilitation in Adults (2014) outline that outcomes of
47 treatment should measure, as a minimum, exercise capacity, dyspnoea and health status and
48 therefore **this service includes completion of the COPD Assessment Test (Jones et al, 2009) and the 6**
49 **minute walk test at the point of initial assessment.**
50
51

52
53 Whilst such standardised assessments assist to meet standards and establish a baseline, the
54 occupational therapy team critically considered that this could promote an initial emphasis on
55 symptoms and disease progression. Therefore, they deliberately also included the Well-being Star
56 (adapted from the Outcomes Star originally developed by Triangle Consulting (Harper, 2004)), and
57 the Hospital Anxiety and Depression (HAD) Scale (Zigmond & Snaith, 1983) in order to more fully
58 understand and measure the biopsychosocial impact of living with lung disease. And whilst
59 recognising a pragmatic need for valid and reliable outcome measurement, the non-standardised
60

1
2
3 initial interview is of equal importance, placing emphasis on a collaborative process to understand a
4 narrative in relation to the person's medical, social and occupational history and facilitating a goal-
5 setting process to help the individual identify what they want to achieve in terms of their own daily
6 activities. At this stage, rather than discussing changes in symptoms or features of the disease (for
7 example a common starting point is 'I want to feel less breathless' or 'I want to be less anxious'),
8 people are encouraged to re-frame and verbalise what they want to change or adjust to in daily life
9 (for example, 'I want to be able to wash and polish my car without needing to stop'). Such goals are
10 revisited regularly within group PR sessions and are integral to the programme as a whole.
11
12
13

14 Although initial assessments are described as an established part of occupational therapy practice
15 across settings (Pentland et al, 2018), the refinement of this assessment process to balance sensitive
16 outcome measurement with a person-centred collaborative process has been an important and
17 deliberate feature and emphasises an alignment with the philosophy of occupational therapy from
18 the outset. Emphasis is given to wellbeing (a concept acknowledged as a state of *being* and *doing*),
19 co-constructing how individuals subjectively interpret their own wellbeing through a narrative, and
20 focussing on person- and occupation-centred goals (Fieldhouse & Bannigan, 2014; Royal College of
21 Occupational Therapists (RCOT), 2015a). Whilst carrying this out in a person's home may feel
22 obvious and intuitive for occupational therapists, situating conversations about occupations within
23 the context in which they occur is another element of deliberate and reflective reasoning.
24
25
26
27
28
29

30 *Flexible and Responsive Patient Education*

31
32 Often PR programmes follow directive models, with pre-determined and professionally-led
33 educational topics or activities delivered in a set order. Yet recent evidence suggests this may
34 undermine people's confidence to self-manage long-term health conditions and instead, positive
35 outcomes may be achieved through personalised and tailored approaches, focussing on
36 conversations with health professionals to discuss goals, actions and identify the elements people
37 require support to manage (Coulter et al, 2015).
38
39
40

41 Self-determination, that is the right of a person to make independent decisions in the process of
42 receiving care, is again emphasised as an important feature of a rehabilitative process (Ekelund,
43 Dahlin-Ivanoff & Eklund, 2013). In line with this, rather than a 'one-size-fits-all', pre-determined
44 programme, this PR model encourages group members to identify the elements of their health and
45 wellbeing where they would value professional support. For pragmatic reasons, some professionally-
46 led sessions are pre-booked, but others involve group choice. **And for all sessions, including those
47 with pre-booked facilitators such as a dietician or respiratory nurse, the education session starts
48 with a group discussion about what they want to know, how this particular topic will help them to
49 manage their own health and wellbeing and involves peer discussion to encourage the group to
50 share support strategies linked to the topic.** Choice within group activities has been supported as a
51 strategy to foster feelings of competency and self-efficacy (Bandura, 1999). Concepts of self-
52 determination and choice can be clearly linked to humanistic perspectives which informed the early
53 development of the occupational therapy profession (Creek, 2014) and which continue to be
54 reflected throughout professional standards and codes of conduct (RCOT, 2015b).
55
56
57
58
59
60

An Enabling Environment for Participation in Occupation

The programme takes place in local gym and community facilities, with groups of people who have self-selected this location and participate in the full 6-week programme together. Not only is this seen as an authentic social and physical environment for the activity of attending the weekly PR session, but it is also potentially an authentic environment for past, current or future occupations. The link to sustainable occupational change is important here and anecdotal feedback suggests many people continue to engage in social activities, or use the same facilities after completion of the formal programme.

The environment in which occupation takes place is understood as a crucial enabler or barrier to optimising performance and engagement and is reflected in models of professional practice (Polatjko, Townsend & Craik, 2007). The design of human and non-human conditions within the environment is linked to the core occupational therapy skill of activity and occupational analysis although the challenge is that a simplified common-sense approach is assumed, when in reality, facilitating environmental conditions to promote engagement in occupation is often complex, non-standardised and difficult to describe (Cole, 2018).

Case Study

James* is a 71 year old man who was recently widowed and has a diagnosis of COPD. James had a stroke 4 years ago which resulted in a left sided weakness and poor balance. He was referred to PR by his Consultant Physician although initially declined to participate, feeling hopeless about anything which required him to go out or do something new.

After a second approach, he consented to the initial assessment and James later acknowledged if he had been required to go out for this initial appointment he would not have gone. Baseline standardised assessments were completed although it was during discussions that he shared feelings of isolation, low mood, anger about his situation and that he was using alcohol heavily to cope. James was asked about his goals at this stage and he could not identify anything he thought he could change. However, James did share that the experience of the initial assessment was a positive one and that sharing his situation with someone who was willing to listen was helpful in itself. Despite James struggling to identify goals related to his everyday life, the occupational therapist shared examples of things she had heard him talk about that had previously held meaning to him and how doing these things again could be goals in the future.

James therefore agreed to start to attend a group programme, choosing a venue already known to him and valuing that the location meant he could stop and go home if he wanted to. At an early stage his occupational therapist was discussing with him how the new occupation of attending the group linked to familiar environments and routines. In early groups, James needed a lot of support to engage in the exercises and, with the occupational therapist mindful of building feelings of self-efficacy, she worked closely with the exercise instructor to grade and adapt the activities and the environment taking in to consideration his left sided weakness, balance and his breathlessness. Socially, he did not say much in the group but discussed later that seeing others making decisions and setting their own goals was a turning point in helping him to see his own situation more

1
2
3 positively. He began to set his own goals within the group, the first of which simply focussed on
4 walking to the gym on his own.
5

6
7 On completion of the programme, repeat standardised assessments evidenced improvements in his
8 activity tolerance and his mood, a reduction in how his COPD was impacting on his overall health
9 and he had also stopped smoking and drinking. He has continued to go to the same gym, attending
10 classes with the same exercise instructor and with the same group. He also bought home exercise
11 equipment to enable him to continue to maintain his positive lifestyle changes.
12
13

14 15 16 **Implications for Practice** 17

18 A synthesis of the features above reveals that rather than being a discreet, or one-off focus on
19 occupation (for example, through a one-off session facilitated by an occupational therapist),
20 occupation is the central feature of this service, underpinned by core knowledge, values and skills of
21 occupational therapy. Reflected in an earlier section, the centrality of exercise and strengthening as
22 a key element of PR programmes has commonly placed physiotherapists in leading roles within PR
23 services, yet conversely it is the centrality of occupation which has strengthened this specific model
24 and resulted in the occupational therapy-led PR programme becoming an established part of the
25 commissioned service provision.
26
27
28

29 Some of the strengths of this service lie in often taken for granted aspects of occupational therapy
30 practice such as an initial assessment in a person's home; a familiar and accessible location for a
31 group; and a collaborative goal-setting process valuing activities which hold personal meaning. It is
32 hoped such reminders will provide important messages for those involved in delivering or
33 developing occupational therapy services within PR. This is particularly pertinent when services may
34 be encouraged to manage costs – for example, by facilitating initial assessments within a maximum
35 time (and therefore perhaps prioritising the standardised elements by default), or undertaking
36 assessments and interventions within clinic environments.
37
38
39

40 However, this synthesis has also revealed some crucial challenges. Firstly, as a profession,
41 occupational therapists struggle to articulate why they are centrally, and perhaps uniquely placed to
42 lead the elements of practice outlined in this analysis. Many professionals have expertise in
43 delivering and facilitating group interventions, and terms such as 'choice', 'person-centredness' and
44 'collaboration' are now part of a more generic discourse. It is not sufficient to use such terms and
45 rely on implicit links to core skills and professional reasoning of occupational therapy but instead
46 occupational therapists must develop vocabulary, grown from a strong theoretical foundation, to
47 describe their contribution.
48
49
50

51 Furthermore, difficulty describing practice— such as initial assessments, goal-setting processes and
52 self-management approaches – can result in these elements becoming varied, non-standardised and
53 difficult to replicate. Whilst this does support flexible and individualised services, it also leads to
54 challenges when delivering a programme across different therapists and locations and difficulty with
55 meaningful evaluation or claims about effectiveness.
56
57
58

59 Noticing such challenges has been an important area of development, and has highlighted future
60 actions for service and professional development and exciting opportunities for research. Looking

1
2
3 ahead, the team are keen to evaluate and compare this PR model through experimental research
4 and recognise it is this which will help to answer important clinical questions about which models
5 and components are most effective. However, research to describe and optimise current service
6 provision (so it can be subsequently implemented and replicated consistently) is a necessary
7 developmental stage before trying to evaluate the effectiveness of a complex intervention (Craig et
8 al, 2008). Qualitative research is therefore planned within this service in order to describe the
9 programme from the perspective of service users and stakeholders and to gain consensus about the
10 best-practice features.
11
12
13
14
15

16 **Key Messages**

17
18 Occupation can be the central tenet of a pulmonary rehabilitation programme with occupational
19 therapists ideally placed to lead and develop services.
20

21
22 However, occupational therapists must clearly articulate the unique knowledge and skills informing
23 occupation-focussed pulmonary rehabilitation programmes in order to inform commissioning of
24 services and to create conditions for robust evaluation and research.
25
26
27

28 **Research Ethics**

29
30 Ethics approval was not required nor sought for this practice analysis. The case study above* is
31 based on a hypothetical case with no identifiable information.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **References**

49 Adeloje D, Chua S, Lee C, Basquill C, Papan A, Theodoratou E, Nair H, Gasevic D, Sridhar D,
50 Campbell H, Yee Chan K, Sheik A and Rudan I (2015) Global and regional estimates of COPD
51 prevalence: systematic review and meta-analysis. *Journal of Global Health*. Available at:
52 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4693508/pdf/jogh-05-020415.pdf> (accessed on 3
53 January 2019)
54

55 Bandura A (1999) Social cognitive theory: an agentic perspective. *Asian Journal of Social Psychology*:
56 2; 21.
57

58
59 British Lung Foundation (2017) Estimating the economic burden of respiratory illness in the UK.
60 Available at: <https://www.blf.org.uk/policy/economic-burden> (accessed on 16 October 2018).

1
2
3 British Thoracic Society (2014) Quality Standards for Pulmonary Rehabilitation in Adults. Available at:
4 [https://www.brit-thoracic.org.uk/document-library/clinical-information/pulmonary-](https://www.brit-thoracic.org.uk/document-library/clinical-information/pulmonary-rehabilitation/bts-quality-standards-for-pulmonary-rehabilitation-in-adults/)
5 [rehabilitation/bts-quality-standards-for-pulmonary-rehabilitation-in-adults/](https://www.brit-thoracic.org.uk/document-library/clinical-information/pulmonary-rehabilitation/bts-quality-standards-for-pulmonary-rehabilitation-in-adults/) (accessed on 3 January
6 2019)
7

8
9 Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I. & Petticrew, M. (2008) Developing and
10 evaluating complex interventions: the new Medical Research Council guidance. *British Medical*
11 *Journal*. Available online at: <https://www.bmj.com/content/bmj/337/bmj.a1655.full.pdf> (accessed
12 on 2 January 2019)
13

14
15 Creek J (2014) The knowledge base of occupational therapy. In Bryant W, Fieldhouse J and Bannigan
16 K (Eds) *Creek's Occupational Therapy and Mental Health – 5th Edition*. Edinburgh: Churchill
17 Livingstone Elsevier.
18

19
20 Cole M (2018) *Group dynamics in occupational therapy: the theoretical basis and practice application*
21 *of group intervention – 5th Edition*. Thorofare, NJ: Slack Incorporated.
22

23
24 Coulter A, Entwistle V, Eccles A, Ryan S, Shepperd S and Perera R (2015) Personalised care planning
25 for adults with chronic or long-term health conditions. *Cochrane Database of Systematic Reviews*.
26 Available at: <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD010523.pub2/full>
27 (accessed on 5th April 2019).
28

29
30 Department of Health (2012) *COPD Commissioning Toolkit: A Resource for Commissioners*. Available
31 at:
32 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/212876/chronic-obstructive-pulmonary-disease-COPD-commissioning-toolkit.pdf)
33 [/212876/chronic-obstructive-pulmonary-disease-COPD-commissioning-toolkit.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/212876/chronic-obstructive-pulmonary-disease-COPD-commissioning-toolkit.pdf) (accessed on 16
34 October 2018).
35

36
37 Ekelund C, Dahlin-Ivanoff S and Eklund K (2013) Self-determination and older people – a concept
38 analysis. *Scandinavian Journal of Occupational Therapy*. Available at:
39 <https://www.tandfonline.com/doi/abs/10.3109/11038128.2013.853832> (accessed on 3 January
40 2019).
41

42
43 Fieldhouse J and Bannigan K (2014) Mental Health and Wellbeing. In Bryant W, Fieldhouse J,
44 Bannigan K (Eds) *Creek's Occupational Therapy and Mental Health – 5th Edition*. Edinburgh: Churchill
45 Livingstone Elsevier.
46

47
48 Global Initiative for Chronic Obstructive Lung Disease (GOLD) (2017) *Global Strategy for the*
49 *Diagnosis, Management and prevention of Chronic Obstructive Pulmonary Disease*. Available at:
50 <https://goldcopd.org> (accessed 16 October 2018).
51

52
53 Harper L (2004) *First research results from the outcome star*. London: St Mungos.
54

55
56 Jones P, Harding G, Berry P, Wiklund I, Chen W and Kline Leidy N (2009) Development and first
57 validation of the COPD Assessment Test. *European Respiratory Journal*. Available at:
58 <https://erj.ersjournals.com/content/erj/34/3/648.full.pdf> (accessed on 3 January 2019).
59

60
61 McCarthy B, Casey D, Devane D, Murphy K, Murphy E, Lacasse Y (2015) Pulmonary rehabilitation for
62 chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*. Available at:
63

1
2
3 <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD003793.pub3/epdf/full> (accessed
4 on 16 October 2018).

5
6 Pentland D, Kantartzis S, Giatsi Clausen M and Witemyre K (2018) *Occupational therapy and*
7 *complexity: defining and describing practice*. London: RCOT.

8
9
10 Polatajko H, Townsend E and Craik J (2007) The Canadian Model of Occupational Performance and
11 Engagement (CMOP-E). In Townsend E and Polatajko H (Eds) *Enabling occupation II: Advancing an*
12 *occupational therapy vision for health, well-being, and justice through occupation*. Ottawa, ON:
13 CAOT Publications.

14
15 Royal College of Occupational Therapists (2015a) *Essential Briefing: Occupation-centred practice*.
16 London: RCOT.

17
18 Royal College of Occupational Therapists (2015b) *Code of Ethics and Professional Conduct*. London:
19 RCOT.

20
21
22 Spruit M, Singh S, Garvey C, ZuWallack R, Nici L, Rochester C, Hill K, Holland A, Lareau S, Man W and
23 Pitta F (2013) An official American Thoracic Society/European Respiratory Society statement: key
24 concepts and advances in pulmonary rehabilitation. *American Journal of Respiratory and Critical*
25 *Care Medicine*. Available at: <https://www.atsjournals.org/doi/abs/10.1164/rccm.201309-1634st>
26 (accessed on 2 January 2019).

27
28
29 Walker BA, Breckner H, Carrier M, Pullen W, Reagan E, Telfer L and Zimmerman T (2016)
30 "Openaccess review to determine best evidence-based practice for COPD". *The Open Journal of*
31 *Occupational Therapy*. Available at: <https://scholarworks.wmich.edu/ojot/vol4/iss2/8/> (accessed on
32 16 October 2018).

33
34
35 Yohannes A and Connolly M (2004) Pulmonary rehabilitation programmes in the UK: a national
36 representative survey. *Clinical Rehabilitation*. Available online at:
37 <https://journals.sagepub.com/doi/abs/10.1191/0269215504cr736oa> (accessed on 3 January 2019).

38
39
40 Zigmond A and Snaith R (1983) The hospital anxiety and depression scale. *Acta Psychiatrica*
41 *Scandinavica*. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6880820> (accessed on 3
42 January 19).

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60