

Northumbria Research Link

Citation: Ratliff, Daniel J. and Bridges, Thomas J. Reduction to modified KdV and its KP-like generalization via phase modulation. *Nonlinearity*, 31 (8). pp. 3794-3813. ISSN 0951-7715

Published by: UNSPECIFIED

URL:

This version was downloaded from Northumbria Research Link: <http://northumbria-test.eprints-hosting.org/id/eprint/54532/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



Northumbria
University
NEWCASTLE



UniversityLibrary

Reduction to modified KdV and its KP-like generalization via phase modulation

Daniel J. Ratliff¹ & Thomas J. Bridges²

1. Department of Mathematical Sciences, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK

2. Department of Mathematics, University of Surrey, Guildford, Surrey, GU2 7XH, UK

E-mail: D.J.Ratliff@lboro.ac.uk, T.Bridges@surrey.ac.uk

20th March 2018

Abstract. The main observation of this paper is that the modified Korteweg-de Vries equation has its natural origin in phase modulation of a basic state such as a periodic travelling wave or more generally a family of relative equilibria. Extension to 2+1 suggests that a modified Kadomtsev-Petviashvili (or a Konopelchenko-Dubrovsky) equation should emerge, but our result shows that there is an additional term which has gone heretofore unnoticed. Thus through the novel application of phase modulation a new equation appears as the 2+1 extension to a previously known one. To demonstrate the theory it is applied to the cubic-quintic Nonlinear Schrödinger (CQNLS) equation, showing that there are relevant parameter values where a modified KP equation bifurcates from periodic travelling wave solutions of the 2+1 CQNLS equation.

PACS numbers: 47.10.Df, 47.35.-i, 42.10, 45.20.Jj

Keywords: nonlinear waves, Lagrangian fields, phase dynamics

Submitted to: *Nonlinearity*

1. Introduction

The modified Korteweg-de Vries equation (mKdV),

$$u_t \pm u^2 u_x + u_{xxx} = 0, \quad (1.1)$$

for the scalar-valued function $u(x, t)$ is most well-known as an integrable member of the KdV family. The interest of this paper will be on how the mKdV (and its 2+1 generalisations)

emerge in a reduction from an arbitrary PDE generated by a Lagrangian. The properties and solutions of the mKdV equation motivate this analysis, but are not the focus of this paper. The mKdV equation is related to the KdV equation through the Miura transformation [14]. However, since the Miura transformation is not bijective, the range of solutions of mKdV is much larger than KdV [25]. For example, the mKdV equation has breather solutions and the cnoidal waves are Benjamin-Feir unstable when (1.1) has the plus sign [8, 25]. The mKdV has been found in a wide range of applications: Alfvén waves, anharmonic lattices, ion acoustic solitons, traffic flow, thin ocean jets, heat pulses in solids, and internal waves (see [25] and references therein).

Typically the KdV or mKdV is derived starting with the dispersion relation in the neighbourhood of a trivial state. The dispersion relation generates the linear terms and an asymptotic expansion or the method of multiples scales is used to generate the nonlinear terms (e.g. [7]). In this paper this strategy is reversed; the basic state is used as the generator of the mKdV equation. This strategy is implemented using *phase modulation*. It is natural since it encapsulates the case of emergence from simple or trivial basic states but also generalises to the case of bifurcation from periodic travelling waves. Phase modulation dramatically enlarges the range of contexts where the mKdV emerges as a model.

The generic system arising from phase modulation is the dispersionless Whitham equations, and at codimension one the KdV equation arises [1], or KP in the case of 2 + 1 with symmetry [18], both associated with long-wave modulation (although the “long-wave” assumption is not explicit, but implicit through the appearance of zero characteristics in generic Whitham theory). In order for the mKdV equation, or its 2 + 1 generalisation, to emerge at least two parameters need to be varied. In 1 + 1, mKdV is one of the natural codimension two modulation equations arising from phase modulation (see Figure 1 below for a classification by codimension).

Phase modulation has its origins in Whitham modulation theory (WMT) [23] but WMT generates dispersionless modulation equations. The idea in phase modulation is to use a more general scaling and a direct ansatz with explicit wavenumber and frequency modulation (e.g. [1, 18, 19, 20]), taking into account natural singularities. With the right scaling dispersion enters naturally. In classical Whitham theory, dispersion can be brought in via amplitude modulation (e.g. §15.5 in [23] and section II of [24]). However, in that theory there is a strong restriction on amplitude, requiring amplitude to be of order ε restricting that theory to the weakly nonlinear case. In phase modulation, dispersion arises naturally with no restriction on amplitude of the basic state.

As in WMT, the starting point is the class of PDEs that are the Euler-Lagrange equation associated with a general Lagrangian density,

$$\mathcal{L}(U) = \int \int \int \mathcal{L}(U, U_t, U_x, U_y) dx dy dt, \quad (1.2)$$

for some vector-valued function $U(x, y, t)$. The theory will be developed in the context of 2 + 1 and this will lead to more than the mKdV equation and indeed more than the modified

Reduction to mKP-KD⁺ via phase modulation 3

KP equation. The theory revolves around the basic state. For definiteness the basic state is assumed to be a periodic travelling wave solution

$$U(x, y, t) = \widehat{U}(kx + my + \omega t + \theta_0) = \widehat{U}(\theta), \quad \widehat{U}(\theta + 2\pi) = \widehat{U}(\theta), \quad (1.3)$$

where (k, m) are the wavenumbers and ω is the frequency, although the theory can be built around any relative equilibrium (cf. [1, 18, 19, 20]). The Lagrangian, averaged over one period, on the wavetrain is then

$$\mathcal{L}(\omega, k, m) = \int_0^{2\pi} \mathcal{L}(\widehat{U}, \omega \widehat{U}_\theta, k \widehat{U}_\theta, m \widehat{U}_\theta) d\theta. \quad (1.4)$$

The Whitham strategy is to substitute $k = \theta_x$, $m = \theta_y$, $\omega = \theta_t$, and take the variation of (1.4) with respect to θ which results in the conservation of wave action,

$$\mathcal{A}(\omega, k, m)_t + \mathcal{B}(\omega, k, m)_x + \mathcal{C}(\omega, k, m)_y = 0, \quad (1.5)$$

where ω , k , and m are considered as functions of x, y, t , and $\mathcal{A} := \mathcal{L}_\omega$, $\mathcal{B} := \mathcal{L}_k$, and $\mathcal{C} := \mathcal{L}_m$. This equation, coupled with the consistency conditions,

$$k_t = \omega_x, \quad m_t = \omega_y, \quad k_y = m_x, \quad (1.6)$$

form the Whitham modulation equations (WMEs). However the four equations (1.5)-(1.6) are a set of *dispersionless* first-order PDEs. To generate the mKdV equation a more general approach is needed.

Our strategy here is to use an explicit ansatz for the modulation. That is the modulation of $\widehat{U}(\theta, \omega, k, m)$ is represented by

$$\widehat{U}(\theta + \varepsilon^a \phi, \omega + \varepsilon^b \Omega, k + \varepsilon^c q, m + \varepsilon^d r) + \varepsilon^e W, \quad (1.7)$$

where W is a remainder, ε is a small parameter, the exponents a, b, c, d, e are problem dependent, and Ω , q , and r depend on slow time and space scales. Singularities in wavenumber-frequency space affect the choice of scales, and are the key mechanism for bringing in dispersion. Substitution of (1.7) into the Euler-Lagrange equation, expansion, and reduction then leads to PDEs for Ω , q and r . The aim is to identify the singularities and scaling that result in a mKdV in $1 + 1$ and its generalisation to $2 + 1$.

The main result of this paper is that when the components $(\mathcal{A}, \mathcal{B}, \mathcal{C})$ of the wave action conservation law, when evaluated on the family of basic states (1.3), have the singularities

$$\mathcal{B}_k = 0, \quad \mathcal{B}_{kk} = 0, \quad \text{and} \quad \mathcal{C}_k = 0, \quad (1.8)$$

in $2 + 1$, then modulation with slow time and space scales

$$X = \varepsilon x, \quad Y = \varepsilon^2 y, \quad T = \varepsilon^3 t, \quad (1.9)$$

Reduction to mKP-KD⁺ via phase modulation 4

and the ansatz

$$\widehat{U}(\theta + \phi, \omega + \varepsilon^3 \Omega, k + \varepsilon q, m + \varepsilon^2 r) + \varepsilon^2 W, \quad (1.10)$$

results in the following modulation equation for q

$$\begin{aligned} \left((\mathcal{A}_k + \mathcal{B}_\omega) q_T + \frac{1}{2} \mathcal{B}_{kkk} q^2 q_X + \mathcal{K} q_{XXX} \right)_X + \mathcal{C}_m q_{YY} \\ + (\mathcal{B}_{km} + \mathcal{C}_{kk})(qq_Y)_X + \mathcal{B}_{km}(q_X \partial_X^{-1} q_Y)_X = 0, \end{aligned} \quad (1.11)$$

with the modulation of Ω and r following from conservation of waves (see (3.3) in §3). This equation will be called mKP-KD⁺ with the plus flagging up the second to last term. In this equation ∂_X^{-1} denotes an antiderivative (when the antiderivative is not well-defined, the term $\partial_X^{-1} q_Y$ is replaced by r). The coefficients in (1.11) are either derivatives of the components of wave action conservation or, in the case of \mathcal{K} , determined by the dispersion relation or a Jordan chain argument, and the details of their construction are given in §3.

Restriction of (1.11) to 1 + 1 gives the mKdV equation in universal form

$$(\mathcal{A}_k + \mathcal{B}_\omega) q_T + \frac{1}{2} \mathcal{B}_{kkk} q^2 q_X + \mathcal{K} q_{XXX} = 0, \quad (1.12)$$

with only the first two conditions in (1.8) required. In 2 + 1 one would expect that the generalisation of (1.12) is mKP

$$\left((\mathcal{A}_k + \mathcal{B}_\omega) q_T + \frac{1}{2} \mathcal{B}_{kkk} q^2 q_X + \mathcal{K} q_{XXX} \right)_X + \mathcal{C}_m q_{YY} = 0. \quad (1.13)$$

Indeed, this is the appropriate generalisation when $\mathcal{B}_{km} = 0$ and this latter condition is satisfied when the basic state and equations are symmetric in y (see §3.1 for precise symmetry condition). When $\mathcal{B}_{km} \neq 0$ the latter two terms and the complete equation in (1.11) are required.

When only one of the last two terms in (1.11) is operational the mKP generalises to,

$$\left((\mathcal{A}_k + \mathcal{B}_\omega) q_T + \frac{1}{2} \mathcal{B}_{kkk} q^2 q_X + \mathcal{K} q_{XXX} \right)_X + \mathcal{C}_m q_{YY} + \mathcal{B}_{km}(q_X \partial_X^{-1} q_Y)_X = 0. \quad (1.14)$$

This form of equation appears in [12, 13], albeit without the representation of the coefficients in terms of the action as here, and in the literature it is called the *Konopelchenko-Dubrovsky equation*, or mKP-KD for short. However, the complete equation (1.11) with the $(qq_Y)_X$ term included is new. With the scaling (1.9) it is natural to include this term.

It is surprising that the additional $(qq_Y)_X$ term in (1.11) has not been noticed before in generalisations of mKdV to 2 + 1. But in most applications the starting basic state is symmetric (for example a 1 + 1 basic state embedded in 2 + 1) and so a direct derivation using an explicit basic state would not see (or need) the non-symmetric terms. It is however the most general form of the generalisation of mKdV to 2 + 1 and an example with symmetry-breaking terms is given in §5 which requires the $(qq_Y)_X$ term.

Reduction to mKP-KD⁺ via phase modulation

5

The term “universal form” is used to indicate that the resulting equation is deduced from the abstract Lagrangian (1.2), and the coefficients are expressed in terms of the components of the action conservation which are defined purely in terms of the Lagrangian, and don't depend on particular PDEs. A key step in relating the properties of conservation laws to the Euler-Lagrange equation is to transform the Lagrangian into multisymplectic form.

The theory is immediately applicable to any PDE generated by a Lagrangian with a family of periodic travelling waves (or more generally a family of relative equilibria). As a consequence, the theory of this paper anticipates the emergence of the mKP-KD⁺ equation from *finite amplitude* waves, rather than just trivial states, in various contexts. An example which demonstrates the application of this theory is given in §4. Starting with the cubic-quintic variant of the Nonlinear Schrödinger equation (CQNLS),

$$i\Psi_t + \Psi_{xx} + \Psi_{yy} - \alpha|\Psi|^2\Psi + i|\Psi|^2(\beta_1\Psi_x + \beta_2\Psi_y) + |\Psi|^4\Psi = 0, \quad (1.15)$$

for the complex-valued function $\Psi(x, y, t)$, and real-valued parameters β_1 , β_2 , and α , with the latter taken to be positive. The symmetric case, $\beta_1 = \beta_2 = 0$, is considered in §4 and it is shown that at particular parameter values along a branch of periodic travelling waves, the equation (1.11) emerges with $\mathcal{B}_{km} = 0$, and after scaling the coefficients, it can be simplified to

$$(\pm q_T + q^2 q_X + q_{XX})_X - q_{YY} = 0. \quad (1.16)$$

In §5 the effect of β_1, β_2 non zero is considered. It is shown that the full generalisation of mKdV to 2 + 1, the mKP-KD⁺ equation (1.11), emerges with all coefficients nonzero.

The derivation of mKdV in 1+1 in this paper completes the classification of codimension two singularities that arise from the breakdown of the KdV equation in phase modulation. In 1 + 1, the KdV equation arises via phase modulation when the WMEs break down at the singularity $\mathcal{B}_k = 0$ and in universal form [1] it is

$$(\mathcal{A}_k + \mathcal{B}_\omega)q_T + \mathcal{B}_{kk}qq_X + \mathcal{K}q_{XX} = 0, \quad (1.17)$$

where $q(X, T)$ is the modulated wavenumber, $X = \varepsilon x$, and $T = \varepsilon^3 t$. There are three ways this equation can degenerate producing codimension two singularities: $\mathcal{A}_k = 0$ or $\mathcal{K} = 0$ or $\mathcal{B}_{kk} = 0$. In each case a new scaling and ansatz produces a different equation. A diagram showing the new codimension two equations that arise when KdV breaks down is in Figure 1. The first case, $\mathcal{A}_k = 0$, is considered in [19] and the result is

$$\mathcal{A}_k = 0 \quad \text{and} \quad \mathcal{B}_{kk} \neq 0, \mathcal{K} \neq 0 \quad \Rightarrow \quad \mathcal{A}_\omega q_{TT} + \left(\frac{1}{2}\mathcal{B}_{kk}q^2 + \mathcal{K}q_{XX}\right)_{XX} = 0,$$

with the additional constraint $\mathcal{A}_\omega \neq 0$ generated, and new scaling. The second case, $\mathcal{K} = 0$, is considered in [16] and the result is

$$\mathcal{K} = 0 \quad \text{and} \quad \mathcal{A}_k \neq 0, \mathcal{B}_{kk} \neq 0 \quad \Rightarrow \quad (\mathcal{A}_k + \mathcal{B}_\omega)q_T + \mathcal{B}_{kk}qq_X + \mathcal{K}_5 q_{XXXX} = 0,$$

with the additional constraint $\mathcal{K}_5 \neq 0$ generated, and new scaling. The third case where $\mathcal{B}_{kk} = 0$ but the other two coefficients in (1.17) are nonzero, and $\mathcal{B}_{kkk} \neq 0$, is considered




Figure 1: The three codimension two modulation equations associated with breakdown of the KdV equation, with the middle bottom case, $\mathcal{B}_{kk} = 0$ considered in this paper.

in this paper. All three of the above have 2 + 1 generalisations as well. The above “classification by codimension” is reminiscent of classifications in bifurcation theory [6]. The above categorization is in the gradient setting and so could potentially be made precise by cataloguing the singularities of the Lagrangian function $\mathcal{L}(\omega, k, m)$ using catastrophe theory or singularity theory for gradient bifurcation problems [2]. However in this case, since the KdV equation (1.17) has only three coefficients, the classification of codimension two singularities is immediate. However more general and higher codimensional singularities in phase dynamics will require a more systematic approach.

The format of this paper is as follows. The abstract properties of the Lagrangian relevant to the modulation theory are presented in §2, including the multisymplectic formulation, linearisation and conservation laws. The phase modulation analysis is undertaken in §3 and the equation (1.11) is derived. The theory is applied to CQNLS in §4. Implications, other directions, and generalisations are discussed in the concluding remarks §6.

2. Lagrangian, conservation laws, and linearisation

The starting point for the theory is PDEs generated by the 2 + 1 abstract Lagrangian (1.2). The theory relies on transforming this density into *multisymplectic form*, and the process to do so has been documented elsewhere (e.g. [1, 19]). This transformation can be interpreted as a multi-dimensional Legendre transform, or simply as a rearrangement of the Lagrangian to isolate the derivatives in each direction. The transformed Lagrangian is

$$\mathcal{L}(Z) = \int \int \int \left[\frac{1}{2} \langle Z, \mathbf{M}Z_t \rangle + \frac{1}{2} \langle Z, \mathbf{J}Z_x \rangle + \frac{1}{2} \langle Z, \mathbf{K}Z_y \rangle - S(Z) \right] dx dy dt,$$

Reduction to mKP-KD⁺ via phase modulation

7

for a new n -component state vector Z , skew symmetric matrices \mathbf{M} , \mathbf{J} and \mathbf{K} and Hamiltonian function S . The first variation of $\mathcal{L}(Z)$, with fixed endpoints on the variations, results in an Euler-Lagrange equation of the form

$$\mathbf{M}Z_t + \mathbf{J}Z_x + \mathbf{K}Z_y = \nabla S(Z), \quad (2.1)$$

where $Z(x, y, t)$ is an n -component state vector. It is assumed that a periodic travelling wave solution, or relative equilibrium solution exists. For definiteness take the basic state to be the former, but the methodology proceeds almost identically for relative equilibrium. The basic state is

$$\widehat{Z}(\theta, \omega, k, m), \quad \theta := kx + my + \omega t + \theta_0, \quad (2.2)$$

where k, m are wavenumbers, ω is a frequency, and θ_0 is a shift along the solution orbit. This solution satisfies

$$(\omega\mathbf{M} + k\mathbf{J} + m\mathbf{K})\widehat{Z}_\theta = \nabla S(\widehat{Z}). \quad (2.3)$$

The linearisation of (2.1) about the basic state defines the linear operator

$$\mathbf{L}V = \mathbf{D}^2 S(\widehat{Z})V - (\omega\mathbf{M} + k\mathbf{J} + m\mathbf{K})V_\theta.$$

Using this definition, and differentiating (2.3), the θ, k, m and ω derivatives of the basic state satisfy

$$\begin{aligned} \mathbf{L}\widehat{Z}_\theta &= 0, & \mathbf{L}\widehat{Z}_k &= \mathbf{J}\widehat{Z}_\theta, \\ \mathbf{L}\widehat{Z}_m &= \mathbf{K}\widehat{Z}_\theta, & \mathbf{L}\widehat{Z}_\omega &= \mathbf{M}\widehat{Z}_\theta. \end{aligned} \quad (2.4)$$

The first of these highlights that $\widehat{Z}_\theta \in \ker(\mathbf{L})$, and for the purposes of this analysis we will assume that the kernel is no larger. As a consequence solvability of inhomogeneous problems involving \mathbf{L} reduces to

$$\mathbf{L}P = Q \quad \text{is solvable when} \quad \langle\langle \widehat{Z}_\theta, Q \rangle\rangle = 0,$$

for an inner product $\langle\langle \cdot, \cdot \rangle\rangle$. For the periodic travelling wave basic state the inner product is

$$\langle\langle P, Q \rangle\rangle := \frac{1}{2\pi} \int_0^{2\pi} \langle P, Q \rangle d\theta,$$

where $\langle \cdot, \cdot \rangle$ is the Euclidean inner product on \mathbb{R}^n .

2.1. Conservation laws

The theory of phase modulation relies on multisymplectic Noether theory, which relates symmetry to conservation laws using the matrices \mathbf{M} , \mathbf{J} and \mathbf{K} . In the present context, where the basic state is a periodic travelling wave, the conservation law of interest is *conservation of wave action* [1, 18, 19]. After evaluating the components of wave action on the basic state and averaging, they are

$$\mathcal{A}(\omega, k, m) = \frac{1}{2} \langle\langle \widehat{Z}, \mathbf{M}\widehat{Z}_\theta \rangle\rangle, \quad \mathcal{B}(\omega, k, m) = \frac{1}{2} \langle\langle \widehat{Z}, \mathbf{J}\widehat{Z}_\theta \rangle\rangle, \quad \mathcal{C}(\omega, k, m) = \frac{1}{2} \langle\langle \widehat{Z}, \mathbf{K}\widehat{Z}_\theta \rangle\rangle. \quad (2.5)$$

Reduction to $mKP-KD^+$ via phase modulation

8

Derivatives of these arise in the theory, and form both necessary emergence conditions and some of the coefficients in the emergent PDE. The derivatives which are needed in the theory are:

$$\begin{aligned}
\mathcal{A}_k &= \langle\langle \widehat{Z}_k, \mathbf{M}\widehat{Z}_\theta \rangle\rangle = \mathcal{B}_\omega, \\
\mathcal{B}_k &= \langle\langle \widehat{Z}_k, \mathbf{J}\widehat{Z}_\theta \rangle\rangle, \\
\mathcal{C}_k &= \langle\langle \widehat{Z}_k, \mathbf{K}\widehat{Z}_\theta \rangle\rangle = \mathcal{B}_m, \\
\mathcal{B}_{kk} &= \langle\langle \widehat{Z}_{kk}, \mathbf{J}\widehat{Z}_\theta \rangle\rangle + \langle\langle \widehat{Z}_k, \mathbf{J}\widehat{Z}_{\theta k} \rangle\rangle, \\
\mathcal{C}_{kk} &= \langle\langle \widehat{Z}_{kk}, \mathbf{K}\widehat{Z}_\theta \rangle\rangle + \langle\langle \widehat{Z}_k, \mathbf{C}\widehat{Z}_{\theta k} \rangle\rangle = \mathcal{B}_{km}, \\
\mathcal{B}_{kkk} &= \langle\langle \widehat{Z}_{kkk}, \mathbf{J}\widehat{Z}_\theta \rangle\rangle + 3\langle\langle \widehat{Z}_{kk}, \mathbf{J}\widehat{Z}_{\theta k} \rangle\rangle.
\end{aligned} \tag{2.6}$$

2.2. Jordan Chain Theory

Aside from the conservation laws, the other main structure which arises within the reduction scheme is that of a Jordan chain. Ultimately, this leads to a characterisation of the dispersion term in the reduced PDE (1.11). The relevant theory needed is as follows. We will be brief as the necessary theory for Jordan chains has been given in our previous papers [1, 18, 19].

The first two equations in (2.4) highlight the presence of a Jordan chain of the form

$$\mathbf{L}\xi_1 = 0, \quad \mathbf{L}\xi_{i+1} = \mathbf{J}\xi_i, \quad i > 0,$$

with $\xi_1 = \widehat{Z}_\theta$, $\xi_2 = \widehat{Z}_k$. Using the notion of solvability introduced earlier, a third element will exist when

$$0 = \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_k \rangle\rangle = -\mathcal{B}_k,$$

by using (2.6). When this derivative vanishes the chain increases in length, allowing for the definition

$$\mathbf{L}\xi_3 = \mathbf{J}\xi_2 = \mathbf{J}\widehat{Z}_k. \tag{2.7}$$

A fourth element is automatic since $\mathbf{L}\xi_2 = \mathbf{J}\xi_3$ is solvable:

$$\langle\langle \widehat{Z}_\theta, \mathbf{J}\xi_3 \rangle\rangle = -\langle\langle \widehat{Z}_k, \mathbf{J}\widehat{Z}_k \rangle\rangle = 0.$$

This paper assumes that the chain terminates at length four, that is; $\mathbf{L}\xi_5 = \mathbf{J}\xi_4$ is not solvable, and so we introduce the quantity

$$\mathcal{H} = \langle\langle \mathbf{J}\widehat{Z}_\theta, \xi_4 \rangle\rangle \neq 0. \tag{2.8}$$

It can be simplified using the Jordan chain

$$\mathcal{H} := \langle\langle \mathbf{J}\widehat{Z}_\theta, \xi_4 \rangle\rangle = \langle\langle \mathbf{L}\xi_2, \xi_4 \rangle\rangle = \langle\langle \xi_2, \mathbf{L}\xi_4 \rangle\rangle = \langle\langle \xi_2, \mathbf{J}\xi_3 \rangle\rangle. \tag{2.9}$$

This quantity will show up as a dispersion coefficient in the phase modulation theory.

Reduction to mKP-KD⁺ via phase modulation 9

3. Reduction to mKP-KD⁺ via phase modulation

The following modulation ansatz is proposed

$$Z(x, y, t) = \widehat{Z}(\theta + \phi, k + \varepsilon q, m + \varepsilon^2 r, \omega + \varepsilon^3 \Omega) + \varepsilon^2 W(\theta + \phi, X, Y, T), \quad (3.1)$$

where ϕ, q, r, Ω are functions of the slow variables (X, Y, T) , and ε , with definitions

$$X = \varepsilon x, \quad Y = \varepsilon^2 y, \quad T = \varepsilon^3 t.$$

That this is the correct ansatz is confirmed *a posteriori*. The modulation frequency and wavenumbers are related by the conservation of waves,

$$\Omega = \phi_T, \quad q = \phi_X, \quad r = \phi_Y, \quad (3.2)$$

which by cross-differentiation give

$$q_T = \Omega_X, \quad r_T = \Omega_Y, \quad r_X = q_Y. \quad (3.3)$$

The idea is to substitute the ansatz (3.1) into (2.1), Taylor expand around $\varepsilon = 0$ and solve the resulting system of equations. The remainder term W is also expanded in an asymptotic series, with

$$W = W_0(\theta + \phi, X, Y, T) + \varepsilon W_1(\theta + \phi, X, Y, T) + \varepsilon^2 W_2(\theta + \phi, X, Y, T) + \dots$$

A key difference from previous analyses of this type is that the remainder term W now depends on ϕ , since the perturbation $\theta \mapsto \theta + \phi$ is of order unity.

The equations at order ε^j , $j = 0, 1, 2, 3, 4$ are needed, with the first three brief, and the mKP-KD⁺ arising at fourth order. The ε^0 equations is just

$$(\omega \mathbf{M} + k \mathbf{J} + m \mathbf{K}) \widehat{Z}_\theta = \nabla S(\widehat{Z}),$$

recovering (2.3) and \widehat{Z} satisfies this equation by assumption. Using the second result of (2.4), the ε^1 equation reduces to

$$(q - \phi_X) \mathbf{L} \widehat{Z}_\theta = 0,$$

recovering the phase consistency condition $q = \phi_X$. Ignoring terms that cancel and using (2.4) again, the ε^2 equation is

$$\mathbf{L} W_0 + (r - \phi_Y) \mathbf{L} \widehat{Z}_m = q_X \mathbf{J} \widehat{Z}_k.$$

The second term on the left vanishes trivially by the phase consistency condition $r = \phi_Y$. For the term on the right hand side, solvability is imposed and generates the condition

$$0 = \langle \langle \widehat{Z}_\theta, \mathbf{J} \widehat{Z}_k \rangle \rangle = -\langle \langle \mathbf{J} \widehat{Z}_\theta, \widehat{Z}_k \rangle \rangle = -\mathcal{B}_k \quad (3.4)$$

1
2
3 *Reduction to mKP-KD⁺ via phase modulation* 10
4

5 This is precisely the condition seen in §2.2 for the chain to be of length 4. Assuming the
6 condition $\mathcal{B}_k = 0$, the equation can be solved for W_0 , giving
7

$$8 \quad W_0 = q_X \xi_3, \quad (3.5)$$

9
10 where ξ_3 has been defined in (2.7), plus an arbitrary amount of homogeneous solution, which
11 drops out in the final analysis.
12

13
14 At third order the equations get a bit more complicated. The ε^3 terms are

$$15 \quad \mathbf{L}W_1 + (\Omega - \phi_T)\mathbf{L}\widehat{Z}_\omega = qq_X(\mathbf{J}\widehat{Z}_{kk} + \mathbf{J}\partial_\theta\xi_3 - \mathbf{D}^3S(\widehat{Z})(\widehat{Z}_k, \xi_3)) \\ 16 \quad + q_Y(\mathbf{K}\widehat{Z}_k + \mathbf{J}\widehat{Z}_m) + q_{XX}\mathbf{J}\xi_3$$

17
18 Since $\Omega = \phi_T$ the second term on the left hand side vanishes. Now, consider the solvability
19 of the right hand side. The q_Y term is solvable when
20

$$21 \quad 0 = \langle\langle \widehat{Z}_\theta, \mathbf{K}\widehat{Z}_k + \mathbf{J}\widehat{Z}_m \rangle\rangle = -\mathcal{B}_m - \mathcal{C}_k.$$

22
23 We assume this condition to be satisfied henceforth in this analysis. The term in q_{XX} is
24 automatically solvable, using the fourth element in the Jordan chain, $\mathbf{L}\xi_4 = \mathbf{J}\xi_3$. The final
25 term we consider is the qq_X term. Assessing solvability,
26

$$27 \quad 0 = \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{kk} \rangle\rangle + \langle\langle \widehat{Z}_\theta, \mathbf{J}\partial_\theta\xi_3 - \mathbf{D}^3S(\widehat{Z})(\widehat{Z}_k, \xi_3) \rangle\rangle \\ 28 \quad = \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{kk} \rangle\rangle + \langle\langle \xi_3, \mathbf{L}\widehat{Z}_{\theta k} \rangle\rangle \quad (3.6) \\ 29 \quad = \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{kk} \rangle\rangle + \langle\langle \widehat{Z}_{\theta k}, \mathbf{J}\widehat{Z}_k \rangle\rangle = -\mathcal{B}_{kk}.$$

30
31 Therefore we require \mathcal{B}_{kk} to vanish for the analysis to continue. This confirms the necessity
32 of all three conditions (1.8) for the emergence of mKP-KD⁺.
33

34
35 When the conditions are met, the solution for W_1 is

$$36 \quad W_1 = qq_X\partial_k\xi_3 + q_{XX}\xi_4 + r_X\zeta, \quad (3.7)$$

37
38 plus an arbitrary amount of homogeneous solution which is neglected as it does not affect
39 the final equation at fourth order. In (3.7) the function ζ is defined by
40

$$41 \quad \mathbf{L}\zeta = \mathbf{K}\widehat{Z}_k + \mathbf{J}\widehat{Z}_m.$$

42
43 The function $\partial_k\xi_3$ in the qq_X term has been deduced by taking the k derivative of the defining
44 equation for ξ_3 ,
45

$$46 \quad \mathbf{D}^3S(\widehat{Z})(\widehat{Z}_k, \xi_3) + \mathbf{D}^2S(\widehat{Z})\partial_k\xi_3 - (\omega\mathbf{M} + k\mathbf{J})\partial_{\theta k}\xi_3 - \mathbf{J}(\xi_3)_\theta = \mathbf{J}\widehat{Z}_{kk},$$

47
48 or

$$49 \quad \mathbf{L}\partial_k\xi_3 = \mathbf{J}\widehat{Z}_{kk} + \mathbf{J}\partial_\theta\xi_3 - \mathbf{D}^3S(\widehat{Z})(\widehat{Z}_k, \xi_3). \quad (3.8)$$

50
51
52
53
54
55
56
57
58
59
60

Reduction to mKP-KD⁺ via phase modulation

11

At fourth order, collecting the necessary terms and simplifying the result,

$$\begin{aligned}
\mathbf{L}W_2 = & q_T(\mathbf{M}\widehat{Z}_k + \mathbf{J}\widehat{Z}_\omega) + q_{XXX}\mathbf{J}\xi_4 + q_{XY}(\mathbf{J}\zeta_1 + \mathbf{K}\xi_3) + r_Y\mathbf{K}\widehat{Z}_m \\
& + qr_X(\mathbf{J}\widehat{Z}_{km} + \mathbf{K}\widehat{Z}_{kk} + \mathbf{J}\zeta_1 - D^3S(\widehat{Z})(\zeta_1, \widehat{Z}_k)) \\
& + r q_X(\mathbf{J}\widehat{Z}_{km} + \mathbf{K}\partial_\theta\xi_3 - D^3S(\widehat{Z})(\widehat{Z}_m, \xi_3)) \\
& + qq_{XX}(D^3S(\widehat{Z})(\widehat{Z}_k, \xi_4) - \mathbf{J}(\xi_4)_\theta - \mathbf{J}\partial_k\xi_3) + (q_X)^2(D^3S(\widehat{Z})(\xi_3, \xi_3) - 2\mathbf{J}\partial_k\xi_3) \\
& + q^2q_X\left(\frac{1}{2}\mathbf{J}\widehat{Z}_{kkk} + \mathbf{J}(\xi)_{\theta k} - \frac{1}{2}D^4S(\widehat{Z})(\widehat{Z}_k, \widehat{Z}_k, \xi_3)\right) \\
& - q^2q_X\left(\frac{1}{2}D^3S(\widehat{Z})(\widehat{Z}_{kk}, \xi_3) + D^3S(\widehat{Z})(\widehat{Z}_k, \partial_k\xi_3)\right).
\end{aligned} \tag{3.9}$$

Solvability of the above is now considered, with the final result being the desired phase equation. Take the inner product of the right hand side with $\widehat{Z}_\theta = \xi_1$. Firstly, the coefficient of the q_T term can be computed to show that

$$\langle\langle \widehat{Z}_\theta, \mathbf{M}\widehat{Z}_k + \mathbf{J}\widehat{Z}_\omega \rangle\rangle = -\mathcal{A}_k - \mathcal{B}_k.$$

Using the definition (2.8), the coefficient of the q_{XXX} term is

$$\langle\langle \widehat{Z}_\theta, \mathbf{J}\xi_4 \rangle\rangle = -\mathcal{H}.$$

Similarly the coefficient of r_Y is

$$\langle\langle \widehat{Z}_\theta, \mathbf{K}\widehat{Z}_m \rangle\rangle = -\mathcal{C}_m.$$

The next considered is the q^2q_X term. Starting from our expression for \mathcal{B}_{kk} ,

$$\begin{aligned}
\mathcal{B}_{kkk} &= \partial_k(\mathcal{B}_{kk}) \\
&= \partial_k\left(-\langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{kk} \rangle\rangle - \langle\langle \widehat{Z}_\theta, \mathbf{J}\partial_\theta\xi_3 \rangle\rangle + \langle\langle \widehat{Z}_\theta, D^3S(\widehat{Z})(\widehat{Z}_k, \xi_3) \rangle\rangle\right) \\
&= -\langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{kkk} + \mathbf{J}\partial_{\theta k}\xi_3 - D^4S(\widehat{Z})(\widehat{Z}_k, \widehat{Z}_k, \xi_3) - D^3S(\widehat{Z})(\widehat{Z}_{kk}, \xi_3) \rangle\rangle \\
&\quad + \langle\langle \widehat{Z}_\theta, D^3S(\widehat{Z})(\widehat{Z}_k, \partial_k\xi_3) \rangle\rangle - \langle\langle \widehat{Z}_{\theta k}, \mathbf{L}\partial_k\xi_3 \rangle\rangle.
\end{aligned}$$

This last term can be manipulated to obtain

$$\langle\langle \widehat{Z}_{\theta k}, \mathbf{L}\partial_k\xi_3 \rangle\rangle = \langle\langle \mathbf{J}\widehat{Z}_{\theta\theta} - D^3S(\widehat{Z})(\widehat{Z}_k, \widehat{Z}_\theta), \partial_k\xi_3 \rangle\rangle = \langle\langle \widehat{Z}_\theta, \mathbf{J}\partial_{\theta k}\xi_3 \rangle\rangle - \langle\langle \widehat{Z}_\theta, D^3S(\widehat{Z})(\widehat{Z}_k, \partial_k\xi_3) \rangle\rangle,$$

and so

$$\mathcal{B}_{kkk} = \langle\langle \widehat{Z}_\theta, D^4S(\widehat{Z})(\widehat{Z}_k, \widehat{Z}_k, \xi_3) + D^3S(\widehat{Z})(\widehat{Z}_{kk}, \xi_3) + 2D^3S(\widehat{Z})(\widehat{Z}_k, \partial_k\xi_3) - \mathbf{J}\widehat{Z}_{kkk} - 2\mathbf{J}\partial_{\theta k}\xi_3 \rangle\rangle. \tag{3.10}$$

This shows that the coefficient of the q^2q_X term is $-\frac{1}{2}\mathcal{B}_{kkk}$.

Reduction to $mKP-KD^+$ via phase modulation

12

Now consider the quadratic terms. Starting with the qq_{XX} term, its coefficient is given by

$$\begin{aligned}
& \langle\langle \widehat{Z}_\theta, D^3 S(\widehat{Z})(\widehat{Z}_k, \xi_4) - \mathbf{J}\partial_\theta \xi_4 - \mathbf{J}\partial_k \xi_3 \rangle\rangle \\
&= \langle\langle \mathbf{J}\widehat{Z}_\theta, \partial_k \xi_3 \rangle\rangle + \langle\langle \mathbf{J}\widehat{Z}_\theta, \partial_\theta \xi_4 \rangle\rangle + \langle\langle \widehat{Z}_k, D^3 S(\widehat{Z})(\widehat{Z}_\theta, \xi_4) \rangle\rangle \\
&= \langle\langle \mathbf{J}\widehat{Z}_\theta, \partial_k \xi_3 \rangle\rangle + \langle\langle \widehat{Z}_k, \mathbf{J}\partial_\theta \xi_3 - D^3 S(\widehat{Z})(\widehat{Z}_\theta, \xi_4) \rangle\rangle + \langle\langle \widehat{Z}_k, D^3 S(\widehat{Z})(\widehat{Z}_\theta, \xi_4) \rangle\rangle \\
&= \langle\langle \mathbf{J}\widehat{Z}_\theta, \partial_k \xi_3 \rangle\rangle + \langle\langle \widehat{Z}_k, \mathbf{J}\partial_\theta \xi_3 \rangle\rangle \\
&= \partial_k \left[\langle\langle \mathbf{J}\widehat{Z}_\theta, \xi_3 \rangle\rangle \right].
\end{aligned} \tag{3.11}$$

In the above, we have used that

$$\mathbf{L}\partial_\theta \xi_4 = \mathbf{J}\partial_\theta \xi_3 - D^3 S(\widehat{Z})(\widehat{Z}_\theta, \xi_4), \tag{3.12}$$

which can be seen by differentiating the defining equation for ξ_4 . The expression inside the bracket is zero since

$$\langle\langle \mathbf{J}\widehat{Z}_\theta, \xi_3 \rangle\rangle = -\langle\langle \widehat{Z}_\theta, \mathbf{J}\xi_3 \rangle\rangle = -\langle\langle \widehat{Z}_\theta, \mathbf{L}\xi_4 \rangle\rangle = -\langle\langle \mathbf{L}\widehat{Z}_\theta, \xi_4 \rangle\rangle = 0. \tag{3.13}$$

The q_X^2 term is shown to vanish in a similar way,

$$\begin{aligned}
& \langle\langle \widehat{Z}_\theta, D^3 S(\widehat{Z})(\xi_3, \xi_3) - 2\mathbf{J}\partial_k \xi_3 \rangle\rangle \\
&= \langle\langle \xi_3, D^3 S(\widehat{Z})(\widehat{Z}_\theta, \xi_3) \rangle\rangle - 2\langle\langle \widehat{Z}_k, \mathbf{J}\partial_\theta \xi_3 \rangle\rangle \\
&= \langle\langle \xi_3, D^3 S(\widehat{Z})(\widehat{Z}_\theta, \xi_3) - 2\mathbf{J}\widehat{Z}_{k\theta} \rangle\rangle \\
&= -\langle\langle \xi_3, \mathbf{L}\partial_\theta \xi_3 + \mathbf{J}\widehat{Z}_{\theta k} \rangle\rangle \\
&= -\langle\langle \mathbf{J}\widehat{Z}_k, \partial_\theta \xi_3 \rangle\rangle - \langle\langle \xi_3, \mathbf{J}\widehat{Z}_{\theta k} \rangle\rangle \\
&= -\langle\langle \mathbf{J}\widehat{Z}_k, \partial_\theta \xi_3 \rangle\rangle + \langle\langle \partial_\theta \xi_3, \mathbf{J}\widehat{Z}_k \rangle\rangle \\
&= 0,
\end{aligned}$$

using results from (3.11) and (3.13) in going from the second to third lines. Therefore both terms, qq_{XX} and q_X^2 , which have the correct scaling but are not conservative, disappear in the solvability condition.

Next consider the mixed terms (with both r and q). Firstly, the qr_X term simplifies to

$$\begin{aligned}
& \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{km} + \mathbf{K}\widehat{Z}_{kk} + \mathbf{J}\partial_\theta \eta - D^3 S(\widehat{Z})(\eta, \widehat{Z}_k) \rangle\rangle \\
&= \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{km} + \mathbf{K}\widehat{Z}_{kk} \rangle\rangle + \langle\langle \eta, \mathbf{J}\widehat{Z}_{\theta\theta} - D^3 S(\widehat{Z})(\widehat{Z}_\theta, \widehat{Z}_k) \rangle\rangle \\
&= \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{km} + \mathbf{K}\widehat{Z}_{kk} \rangle\rangle + \langle\langle \mathbf{J}\widehat{Z}_m + \mathbf{K}\widehat{Z}_k, \widehat{Z}_{\theta k} \rangle\rangle \\
&= -(\langle\langle \widehat{Z}, \mathbf{J}\widehat{Z}_{\theta km} \rangle\rangle + \langle\langle \widehat{Z}_m, \mathbf{J}\widehat{Z}_{\theta k} \rangle\rangle) - (\langle\langle \widehat{Z}, \mathbf{K}\widehat{Z}_{\theta kk} \rangle\rangle + \langle\langle \widehat{Z}_k, \mathbf{K}\widehat{Z}_{\theta k} \rangle\rangle) \\
&= -\mathcal{B}_{km} - \mathcal{C}_{kk} \quad (\text{noting that } \mathcal{B}_{km} = \mathcal{C}_{kk}).
\end{aligned}$$

Reduction to mKP-KD⁺ via phase modulation

13

Similarly, for the rq_X term,

$$\begin{aligned}
 & \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{km} + \mathbf{K}\partial_\theta\xi_3 - D^3S(\widehat{Z})(\xi_3, \widehat{Z}_m) \rangle\rangle \\
 &= \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{km} \rangle\rangle + \langle\langle \xi_3, \mathbf{K}\widehat{Z}_{\theta\theta} - D^3S(\widehat{Z})(\widehat{Z}_\theta, \widehat{Z}_m) \rangle\rangle \\
 &= \langle\langle \widehat{Z}_\theta, \mathbf{J}\widehat{Z}_{km} \rangle\rangle + \langle\langle \mathbf{J}\widehat{Z}_k, \widehat{Z}_{\theta m} \rangle\rangle \\
 &= -(\langle\langle \widehat{Z}, \mathbf{J}\widehat{Z}_{\theta km} \rangle\rangle + \langle\langle \widehat{Z}_m, \mathbf{J}\widehat{Z}_{\theta k} \rangle\rangle) \\
 &= -\mathcal{B}_{km}.
 \end{aligned}$$

Combining the results of the above calculations, we have shown that the equation for W_2 in (3.9) is solvable when q, r satisfy the PDE

$$(\mathcal{A}_k + \mathcal{B}_\omega)q_T + \frac{1}{2}\mathcal{B}_{kkk}q^2q_X + \mathcal{K}q_{XXX} + \mathcal{C}_m r_Y + (\mathcal{B}_{km} + \mathcal{C}_{kk})qr_X + \mathcal{B}_{km}rq_X = 0. \quad (3.14)$$

Differentiating with respect to X and introducing the antiderivative to solve $r_X = q_Y$ for r recovers (1.11). Alternately, expressing $q = \phi_X$ and $r = \phi_Y$ in terms of the phase in 3.14), the pure phase version of (1.11) is

$$\left((\mathcal{A}_k + \mathcal{B}_\omega)\phi_T + \frac{1}{6}\mathcal{B}_{kkk}\phi_X^3 + \mathcal{K}\phi_{XXX} \right)_X + \mathcal{C}_m\phi_{YY} + (\mathcal{B}_{km} + \mathcal{C}_{kk})\phi_X\phi_{XY} + \mathcal{B}_{km}\phi_Y\phi_{XX} = 0. \quad (3.15)$$

3.1. Reflection symmetric basic state

If the equation has a reflection symmetry in the y -direction and the basic state inherits this symmetry, it is shown in §7 of [18] that the third component of the wave action conservation law in (2.5) satisfies

$$\mathcal{C}(k, -m, \omega) = -\mathcal{C}(\omega, k, m).$$

With this symmetry it is immediate that

$$\mathcal{C}_k \Big|_{m=0} = \mathcal{C}_{kk} \Big|_{m=0} = 0. \quad (3.16)$$

Hence when the basic state is aligned with the x -axis ($m = 0$) the third necessary condition in (1.8) is automatically satisfied, and the coefficients in the latter two terms in (1.11) vanish simplifying the emergent phase equations to the mKP equation

$$\left((\mathcal{A}_k + \mathcal{B}_\omega)q_T + \frac{1}{2}\mathcal{B}_{kkk}q^2q_X + \mathcal{K}q_{XXX} \right)_X + \mathcal{C}_mq_{YY} = 0, \quad (3.17)$$

with the only necessary conditions being $\mathcal{B}_k = \mathcal{B}_{kk} = 0$.

Indeed, in applications it is very often the case that the basic state is symmetric about the x -axis so that $\mathcal{C}_k = \mathcal{C}_{kk} = 0$ automatically. Another point of view is to note that this symmetric case can be understood by noting that then $\mathcal{L}(\omega, k, m) = \widehat{\mathcal{L}}(\omega, \kappa)$ where $\kappa = \sqrt{k^2 + m^2}$ and in particular is an even function of both k and m . This observation leads directly to the symmetry expressions (3.16).

Reduction to $mKP-KD^+$ via phase modulation 14

4. Example: reduction of CQNLS to mKP

The cubic-quintic NLS (CQNLS) equation in standard form is given in (1.15). It appears in the optics literature, describing the propagation of light pulses [9], and in the literature on Bose-Einstein condensates [11]. A self-consistent derivation of CQNLS is given [15]. This example is used here as it is the simplest example with the singularities $\mathcal{B}_k = \mathcal{B}_{kk} = 0$ in the family of travelling waves and it can be explicitly analyzed in some completeness.

The CQNLS equation is the Euler-Lagrange equation associated with the Lagrangian

$$\mathcal{L} = \int_{t_1}^{t_2} \int_{x_1}^{x_2} \left[\frac{1}{2}i(\bar{\Psi}\Psi_t - \Psi\bar{\Psi}_t) - |\Psi_x|^2 - |\Psi_y|^2 - \frac{1}{2}\alpha|\Psi|^4 + \frac{1}{3}|\Psi|^6 \right] dxdt, \quad (4.1)$$

for real constant α , leading to the equation

$$i\Psi_t + \nabla^2\Psi - \alpha|\Psi|^2\Psi + |\Psi|^4\Psi = 0. \quad (4.2)$$

The CQNLS equation (4.2) has a three-parameter family of exact periodic travelling wave solutions

$$\Psi(x, y, t) = \Psi_0 e^{i\theta}, \quad \theta = kx + my + \omega t. \quad (4.3)$$

Substitution of this form of solution into (4.2) gives the amplitude as a function of the wavenumber and frequency,

$$\Delta := |\Psi_0|^4 - \alpha|\Psi_0|^2 - \omega - k^2 - m^2 = 0. \quad (4.4)$$

Rearranging, $\Delta = 0$ is equivalent to

$$\left(|\Psi_0|^2 - \frac{1}{2}\alpha \right)^2 - k^2 = \omega + m^2 + \frac{1}{4}\alpha^2, \quad (4.5)$$

giving the necessary conditions for existence

$$|\Psi_0|^2 > 0 \quad \text{and} \quad \omega + k^2 + m^2 + \frac{1}{4}\alpha^2 > 0.$$

In the $(k, |\Psi_0|)$ plane the set $\Delta^{-1}(0)$ is a family of hyperbola parameterised for each fixed (m, α) by ω .

The class of solutions (4.3) can also be considered as a relative equilibrium associated with the $SO(2)$ symmetry; that is, $e^{is}\Psi(x, y, t)$ is a solution of (4.2) whenever $\Psi(x, y, t)$ is a solution, for any $s \in \mathbb{R}$. The conservation law for this symmetry is $A_t + B_x + C_y = 0$ with

$$A = \frac{1}{2}|\Psi|^2, \quad B = \frac{1}{2}(\Psi^*\Psi_x - \Psi_x^*\Psi), \quad C = \frac{1}{2}(\Psi^*\Psi_y - \Psi_y^*\Psi), \quad (4.6)$$

where $*$ denotes complex conjugation. This conservation law can be confirmed using Noether theory or by direct calculation. It is independent of the nonlinearity and so it is the same as the conservation law for the 2+1 cubic NLS equation. In this latter case, this conservation law

Reduction to mKP-KD⁺ via phase modulation

15

is known to represent mass conservation in the theory of water waves and power conservation in optics applications.

Evaluating the components of the conservation law (4.6) on the family of basic states gives

$$\mathcal{A} = \frac{1}{2}|\Psi_0|^2, \quad \mathcal{B} = k|\Psi_0|^2, \quad \mathcal{C} = m|\Psi_0|^2, \quad (4.7)$$

with $|\Psi_0|^2$ expressing using (4.4) as a function of (ω, k, m) . In looking for singular points, in the family of periodic travelling waves, where mKdV emerges, one observation is immediate: $\mathcal{C}(\omega, k, m)$ is an odd function of m , since $|\Psi_0|^2$ is an even function of m , and so

$$\mathcal{C}_k \Big|_{m=0} = \mathcal{C}_{kk} \Big|_{m=0} = 0.$$

Therefore, set $m = 0$, and then we need only identify values of (ω, k) where $\mathcal{B}_k = 0$ and $\mathcal{B}_{kk} = 0$, and the emergent equation will be mKP rather than mKP-KD or mKP-KD⁺ due to the fact that $\mathcal{C}_{kk} = \mathcal{B}_{km} = 0$.

Differentiating \mathcal{B} twice gives

$$\mathcal{B}_k = u + ku_k \quad \text{and} \quad \mathcal{B}_{kk} = 2u_k + ku_{kk}, \quad \text{with } u := |\Psi_0|^2,$$

and by also differentiating (4.5) twice,

$$(2u - \alpha)u_k = 2k \quad \text{and} \quad (2u - \alpha)u_{kk} + 2u_k^2 = 2.$$

Combining the existence condition (4.4), the condition $\mathcal{B}_k = 0$ and the condition $\mathcal{B}_{kk} = 0$, and setting $m = 0$, leads to the three simultaneous nonlinear equations. These equations can be best visualised as three curves in the (u, k, ω) plane,

$$\begin{aligned} \Delta = 0 & \quad \left(u - \frac{1}{2}\alpha\right)^2 - k^2 = \omega + \frac{1}{4}\alpha^2 \\ \mathcal{B}_k = 0 & \quad \left(u - \frac{1}{4}\alpha\right)^2 + k^2 = \frac{1}{16}\alpha^2 \\ \mathcal{B}_{kk} = 0 & \quad \left(u - \frac{1}{2}\alpha\right)^2 - \frac{1}{3}k^2 = 0. \end{aligned} \quad (4.8)$$

The existence set $\Delta^{-1}(0)$ is a hyperbola, the $\mathcal{B}_k = 0$ curve is a circle, and the $\mathcal{B}_{kk} = 0$ curve consists of two straight lines. Where the three surfaces intersect correspond to points where the mKP-KD⁺ equation arises, occurring when the hyperbola is tangent to the circle, with an example of this given in figure 2. The intersection occurs when

$$\omega = -\frac{9}{32}\alpha^2, \quad u := |\Psi_0|^2 = \frac{3}{8}\alpha, \quad k^2 = \frac{3}{64}\alpha^2, \quad (4.9)$$

with α any positive number. Note that there are two distinct k values due to the symmetry with respect to k in (4.4).

This calculation confirms that there exist parameter values where the mKP can potentially emerge. Now compute the values of the coefficients in (1.11), evaluated at the singular point (4.9)

$$\mathcal{A}_k = -4\frac{k}{\alpha}, \quad \mathcal{B}_{kkk} = -\frac{96}{\alpha}, \quad \mathcal{B}_{km} = 0, \quad \mathcal{C}_m = |\Psi_0|^2 = \frac{3}{8}\alpha = \frac{8}{\alpha}k^2. \quad (4.10)$$




Figure 2: An example of the surfaces defined in (4.8) for $\alpha = 4$. The surfaces corresponding to these conditions are coloured green, grey and orange respectively. The intersections between the second and third conditions with the first are highlighted by the black and red lines respectively. The second image shows the intersection between these lines from above and thus the points where the mKP-KD⁺ equation emerges.

In summary, just knowing that CQNLS comes from a Lagrangian and has a family of periodic travelling waves, with the symmetry in m and the singularities $\mathcal{B}_k = 0$ and $\mathcal{B}_{kk} = 0$, it is immediate that there are parameter values where the mKP equation emerges in the form

$$\left(-8\frac{k}{\alpha} q_T - \frac{48}{\alpha} q^2 q_X + \mathcal{K} q_{XXX} \right)_X + \frac{8}{\alpha} k^2 q_{YY} = 0. \quad (4.11)$$

It remains to compute the X -dispersion coefficient \mathcal{K} . This coefficient can be obtained by using the dispersion relation of (4.2) linearised about the family of basic states, or it can be calculated using the Jordan chain argument given in §2.2. The calculation is given below using the Jordan chain argument, and it is found that

$$\mathcal{K} = -\frac{2}{\alpha}. \quad (4.12)$$

Substitution into (4.11) and simplifying gives the final form for the mKP emerging from CQNLS at the singular point (4.9),

$$\left(k q_T + 6 q^2 q_X + \frac{1}{4} q_{XXX} \right)_X - k^2 q_{YY} = 0, \quad (4.13)$$

noting that $k = \pm\alpha\sqrt{3}/8$. By scaling it can be put in the canonical form (1.16). The resulting mKP equation is focussing and so already in the 1 + 1 case the periodic travelling

Reduction to mKP-KD⁺ via phase modulation

17

waves can be Benjamin-Feir unstable [8]. In the 2 + 1 case it is modified KP-I and so the solitary wave solutions will be transverse unstable.

4.1. The dispersion coefficient \mathcal{K}

To confirm the expression (4.12) for the dispersion coefficient the Jordan chain argument in §2.2 is used. Since \mathcal{K} is associated with X -dispersion, it can be calculated using the 1D version of CQNLS,

$$i\Psi_t + \Psi_{xx} - \alpha|\Psi|^2\Psi + |\Psi|^4\Psi = 0. \quad (4.14)$$

First set up the multisymplectic structure \mathbf{M} , \mathbf{J} and S , and construct the linearisation about the family of basic states. Let $\Psi = a_1 + ia_2$ and introduce $\mathbf{a} = (a_1, a_2)$ and $\mathbf{b} = \mathbf{a}_x$. Then the multisymplectic formulation of (4.14) is

$$\mathbf{M}Z_t + \mathbf{K}Z_x = \nabla S(Z), \quad (4.15)$$

where,

$$Z = \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix}, \quad \mathbf{M} = \begin{pmatrix} -\boldsymbol{\sigma} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix}, \quad \mathbf{J} = \begin{pmatrix} \mathbf{0} & -\mathbf{I} \\ \mathbf{I} & \mathbf{0} \end{pmatrix}, \quad \boldsymbol{\sigma} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

and

$$S(Z) = \frac{1}{2}\mathbf{b} \cdot \mathbf{b} - \frac{\alpha}{4}(\mathbf{a} \cdot \mathbf{a})^2 + \frac{1}{6}(\mathbf{a} \cdot \mathbf{a})^3.$$

Just a summary of the calculations are given since analysis of this type appears in previous papers [1, 18, 19]. The basic state in the multisymplectic representation is

$$\widehat{Z}(\theta, k, m, \omega) = G_\theta \begin{pmatrix} \widehat{\mathbf{a}} \\ \widehat{\mathbf{b}} \end{pmatrix},$$

with $|\widehat{\mathbf{a}}|^4 - \alpha|\widehat{\mathbf{a}}|^2 - \omega - k^2 = 0$, $\widehat{\mathbf{b}} = k\boldsymbol{\sigma}\widehat{\mathbf{a}}$, and

$$G_\theta := R_\theta \oplus R_\theta, \quad R_\theta := \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

Linearising (4.15) about the family of basic states generates the linear operator

$$\mathbf{L} := D^2S(\widehat{Z}_\theta) - k\mathbf{J}\partial_\theta - \omega\mathbf{M}\partial_\theta,$$

or, after substitution,

$$\mathbf{L} := \begin{bmatrix} (4|\mathbf{a}|^2 - 2\alpha)\mathbf{a}\mathbf{a}^T + k^2\mathbf{I} & k\boldsymbol{\sigma} \\ -k\boldsymbol{\sigma} & \mathbf{I} \end{bmatrix}.$$

The kernel of \mathbf{L} is spanned by

$$\xi_1 := \widehat{Z}_\theta = G_\theta \begin{pmatrix} \boldsymbol{\sigma}\widehat{\mathbf{a}} \\ -k\widehat{\mathbf{a}} \end{pmatrix}.$$

Reduction to mKP-KD⁺ via phase modulation

18

The next two members of the Jordan chain satisfy $\mathbf{L}\xi_2 = \mathbf{J}\xi_1$ and $\mathbf{L}\xi_3 = \mathbf{J}\xi_2$. These equations are easily solved explicitly giving

$$\xi_2 = G_\theta \begin{pmatrix} \nu \hat{\mathbf{a}} \\ (1 + k\nu)\sigma \hat{\mathbf{a}} \end{pmatrix} \quad \text{and} \quad \xi_3 = G_\theta \begin{pmatrix} 0 \\ \nu \hat{\mathbf{a}} \end{pmatrix},$$

with

$$\nu := \frac{k}{(2|\hat{\mathbf{a}}|^2 - \alpha)|\hat{\mathbf{a}}|^2}.$$

Now using the formula (2.9),

$$\mathcal{K} = \langle\langle \xi_2, \mathbf{J}\xi_3 \rangle\rangle = -\nu^2 |\hat{\mathbf{a}}|^2 = -\frac{2}{\alpha}.$$

Confirming (4.12).

5. Example: breaking transverse symmetry and mKP-KD⁺

Now consider the CQNLS with the β -symmetry breaking terms (1.15). It is sufficient to take $\beta_1 = \beta_2 = \beta$ in which case the starting point for the analysis is

$$i\Psi_t + \nabla^2 \Psi - \alpha |\Psi|^2 \Psi + i\beta |\Psi|^2 (\Psi_x + \Psi_y) + |\Psi|^4 \Psi = 0, \quad (5.1)$$

for complex amplitude $\Psi(x, y, t)$ and real constants $\alpha > 0$ and β . The form of this equation is motivated by the analysis of TUSZYŃSKI & DIXON [21] (see equation (29) in that paper) and the analysis of TUSZYŃSKI & KIBLER [22] (see equation (26) in that paper), although here the coefficients are chosen so that it is Lagrangian generated. In 1 + 1 this equation arises in the theory of water-waves near the critical value $kh \approx 1.363$ for modulation of finite depth Stokes waves (cf. JOHNSON [10]).

The basic state is the same as the previous section, $\Psi(x, y, t) = \Psi_0 e^{i\theta}$, but with a new amplitude equation,

$$\Delta := |\Psi_0|^4 - (\alpha + \beta(k + m))|\Psi_0|^2 - \omega - (k^2 + m^2) = 0. \quad (5.2)$$

The components of the conservation law along this family of solutions, generalising (4.7), are

$$\mathcal{A} = \frac{1}{2} |\Psi_0|^2, \quad \mathcal{B} = k |\Psi_0|^2 + \frac{\beta}{4} |\Psi_0|^4, \quad \mathcal{C} = m |\Psi_0|^2 + \frac{\beta}{4} |\Psi_0|^4.$$

To lighten the notation define $u = |\Psi_0|^2$, then the conservation laws are

$$\mathcal{A}(\omega, k, m) = \frac{1}{2} u, \quad \mathcal{B}(\omega, k, m) = ku + \frac{1}{4} \beta u^2, \quad \text{and} \quad \mathcal{C}(\omega, k, m) = mu + \frac{1}{4} \beta u^2,$$

with u determined as a function of (ω, k, m) from the quadratic

$$\Delta := u^2 - (\alpha + \beta(k + m))u - \omega - (k^2 + m^2) = 0.$$

Reduction to mKP-KD⁺ via phase modulation

19

The calculations proceed as in §4, although the calculations are a bit more lengthy and MAPLE is used; hence just a summary is given. It is necessary to solve the three equations $\mathcal{B}_k = \mathcal{B}_{kk} = \mathcal{C}_k = 0$. Explicit expressions can be obtained but it is easiest to present them in terms of the amplitude at criticality.

First of all, the criticality in \mathcal{C} may be written as

$$\mathcal{C}_k = \left(m + \frac{\beta}{2}u\right)u_k = 0. \quad (5.3)$$

Choosing $u_k = 0$ leads to the solution becoming trivial when the other constraints are imposed, and so we must choose $m = -\frac{\beta}{2}u \neq 0$ in order for this condition to be satisfied. Thus, the role of the transverse symmetry breaking terms becomes apparent, since now a nontrivial m is required for the analysis to proceed. Next, note that the criticality conditions in \mathcal{B} can be written as

$$\begin{aligned} \mathcal{B}_k &= \left(k + \frac{\beta}{2}u\right)u_k + u = 0 \quad \implies \quad u_k = -\frac{u}{k + \frac{\beta}{2}u}, \\ \mathcal{B}_{kk} &= \left(k + \frac{\beta}{2}u\right)u_{kk} + 2u_k + \frac{\beta}{2}u_k^2 = 0 \quad \implies \quad u_{kk} = \frac{u(2k + \frac{\beta}{2}u)}{\left(k + \frac{\beta}{2}u\right)^3}. \end{aligned} \quad (5.4)$$

Before we may write down \mathcal{B}_{kkk} , we need u_{kkk} , which may be found by differentiating the defining expression for u ,

$$u^2 - (\alpha + \beta(k + m))u - \omega - k^2 - m^2 = 0,$$

three times. This gives:

$$\begin{aligned} 2u_{kkk} \left(u - \frac{\alpha + \beta(k + m)}{2}\right) + 6u_k u_{kk} - 3\beta u_{kk} &= 0, \\ \implies u_{kkk} = \frac{3\beta - 6u_k}{2\left(\left(\frac{\beta^2}{4} + 1\right)u - \frac{\alpha + \beta k}{2}\right)} u_{kk} &= \frac{3}{2} \frac{\beta\left(k + \frac{\beta}{2}u\right) + 2u}{\left(\left(\frac{\beta^2}{4} + 1\right)u - \frac{\alpha + \beta k}{2}\right)} \frac{u(2k + \frac{\beta}{2}u)}{\left(k + \frac{\beta}{2}u\right)^4}, \end{aligned} \quad (5.5)$$

using (5.4). Thus, we find that

$$\begin{aligned} \mathcal{B}_{kkk} &= \left(k + \frac{\beta}{2}u\right)u_{kkk} + 3u_{kk} + \frac{3\beta}{2}u_k u_{kk} \\ &= \frac{3u(2k + \frac{\beta}{2}u)}{2\left(k + \frac{\beta}{2}u\right)^4} \left(2k + \frac{(\beta\left(k + \frac{\beta}{2}u\right) + 2u)\left(k + \frac{\beta}{2}u\right)}{\left(\left(\frac{\beta^2}{4} + 1\right)u - \frac{\alpha + \beta k}{2}\right)}\right). \end{aligned}$$

Similarly, we find that

$$\mathcal{A}_k = \frac{1}{2}u_k = -\frac{u}{2\left(k + \frac{\beta}{2}u\right)}, \quad \mathcal{E}_{kk} = \frac{\beta}{2}u_k^2 = \frac{\beta u^2}{2\left(k + \frac{\beta}{2}u\right)^2}, \quad \mathcal{E}_m = u.$$

Reduction to mKP-KD⁺ via phase modulation

20

The Jordan chain analysis is only slightly altered, giving

$$\mathcal{K} = -\frac{u_k^2}{4u} = -\frac{u}{4\left(k + \frac{\beta}{2}u\right)^2}.$$

Therefore, the mKP-KD⁺ is (after multiplying by $k + \frac{\beta}{2}u$ which is nonzero)

$$\left[a_1 q_T + a_2 q^2 q_X + a_3 q_{XX} \right]_X + a_4 q_{YY} + a_5 \left[2(qq_Y)_X + (q_X \partial_X^{-1} q_Y)_X \right] = 0,$$

with

$$\begin{aligned} a_1 &= -u \\ a_2 &= \frac{3u\left(2k + \frac{\beta}{2}u\right)}{4\left(k + \frac{\beta}{2}u\right)^3} \left(2k + \frac{\left(\beta\left(k + \frac{\beta}{2}u\right) + 2u\right)\left(k + \frac{\beta}{2}u\right)}{\left(\frac{\beta^2}{4} + 1\right)u - \frac{\alpha + \beta k}{2}} \right) \\ a_3 &= -\frac{u}{4\left(k + \frac{\beta}{2}u\right)} \\ a_4 &= u\left(k + \frac{\beta}{2}u\right) \\ a_5 &= \frac{\beta u^2}{2\left(k + \frac{\beta}{2}u\right)}. \end{aligned}$$

This equation is in some ways more complicated than the original CQNLS. However, a wide range of solutions can be more easily obtained, and in general one will not always be starting with a simple equation.

6. Concluding remarks

In this paper the x -direction was preferred in that it was the singularities $\mathcal{B}_k = \mathcal{B}_{kk} = 0$ that underpinned the derivation. There is an alternative theory where time is the preferred direction. In the 1 + 1 case the singularities (1.8) are replaced by $\mathcal{A}_\omega = \mathcal{A}_{\omega\omega} = 0$ and the resulting modulation equation is

$$\left(\mathcal{B}_\omega + \mathcal{A}_k\right)\Omega_X + \frac{1}{2}\mathcal{A}_{\omega\omega\omega}\Omega^2\Omega_T + \mathcal{M}\Omega_{TT} = 0, \quad (6.1)$$

with \mathcal{M} determined by a \mathbf{M} -based Jordan chain. This type of equation (with x and t reversed) arises when the spatial development of a wave is of interest and it has been studied when the nonlinearity is quadratic (e.g. equation (8) in [3]) where it is called the KdV equation in signalling coordinates. Hence (6.1) is the mKdV equation in signalling coordinates. The equation (6.1) has 2 + 1 generalisations, and could also be oriented so that y is the preferred direction.

Although related to the KdV equation, the mKdV equation has different structure and a wider range of solutions. For example the focusing mKdV equation has periodic travelling wave solutions which are Benjamin-Feir unstable [8]. It is shown in [4, 5] that this Benjamin-Feir instability can be the catalyst for an energy cascade leading to a continuous spectrum and a highly-complex wave field.

Reduction to mKP-KD⁺ via phase modulation

21

When \mathcal{B}_{kk} is not zero but small, for example $\mathcal{B}_{kk} = \varepsilon\mu$, the mKP-KD⁺ has an unfolding into the 2 + 1 Gardner system which includes both cubic and quadratic nonlinearities.

$$\begin{aligned} & \left((\mathcal{A}_k + \mathcal{B}_\omega)q_T + \mu qq_X + \frac{1}{2}\mathcal{B}_{kkk}q^2q_X + \mathcal{K}q_{XXX} \right)_X \\ & + \mathcal{C}_m q_{YY} + (\mathcal{B}_{km} + \mathcal{C}_{kk})(qq_Y)_X + \mathcal{B}_{km}(q_X \partial_X^{-1} q_Y)_X = 0. \end{aligned}$$

The most important potential generalisation of the theory in this paper is to the case where the symmetry has dimension greater than one, the case of *multiphase modulation*. The case of multiphase modulation in the case of a codimension one singularity leading to KdV is developed in [20]. The generalisation of this theory to the codimension two case is of great interest since the most well-known example of the emergence of the mKdV equation is in multi-layer stratified shallow water hydrodynamics [7] and that case can be viewed as multiphase modulation with a multi-dimensional affine symmetry group. Some progress in this direction is in [17], where a theory for codimension two multiphase modulation leading the Boussinesq equation (a multiphase generalisation of the codimension two singularity in the left of Figure 1) is presented.

Acknowledgements

The first author was funded by an EPSRC PhD studentship with grant number EP/L505092/1.

- [1] BRIDGES T 2013 A universal form for the emergence of the Korteweg–de Vries equation, *Proc. Roy. Soc. Lond. A* **469** 20120707.
- [2] BRIDGES T & FURTER J 1993 *Singularity Theory and Equivariant Symplectic Maps*, Lect. Notes Math. **1558**, (Berlin: Springer-Verlag).
- [3] CAPUTO J AND STEPANYANTS Y 2003 Bore formation, evolution and disintegration into solitons in shallow inhomogeneous channels, *Nonl. Proc. Geophys.* **10** 407–424.
- [4] DUTYKH D AND TOBISCH E 2015 Observation of the inverse energy cascade in the modified Korteweg–de Vries equation, *Euro. Phys. Lett.* **107** 14001.
- [5] DUTYKH D AND TOBISCH E 2015 Direct dynamical energy cascade in the modified KdV equation, *Physica D* **297** 76–87.
- [6] GOLUBITSKY M, STEWART I, & SCHAEFFER D 1988 *Singularities and Groups in Bifurcation Theory, Volume II*, (New York: Springer-Verlag).
- [7] GRIMSHAW R, PELINOVSKY E, TALIPOVA T 1997 The modified Korteweg–de Vries equation in the theory of large-amplitude internal waves, *Nonl. Proc. Geophys.* **4** 237–250.
- [8] GRIMSHAW R, PELINOVSKY D, PELINOVSKY E AND TALIPOVA, T 2001 Wave group dynamics in weakly nonlinear long-wave models, *Physica D* **159** 35–57.
- [9] HERRMANN J 1992 Bistable bright solitons in dispersive media with a linear and quadratic intensity-dependent refraction index change, *Opt. Commun.* **87**, 161–165.
- [10] JOHNSON, R S 1977 On the modulation of water waves in the neighbourhood of $kh \approx 1.363$, *Proc. R. Soc. Lond. A* **357** 131–141.
- [11] KEVREKIDIS P, FRANTZESKAKIS D, & CARRETERO-GONZÁLEZ R 2015 *The Defocusing Nonlinear Schrödinger Equation*, (Philadelphia: SIAM).

- [12] KONOPELCHENKO B On the gauge-invariant description of the evolution equations integrable by Gelfand-Dikij spectral problems, *Phys. Lett. A* **92** 323–327.
- [13] KONOPELCHENKO B AND DUBROVSKY V 1984 Some new integrable nonlinear evolution equations in 2+1 dimensions, *Phys. Lett. A* **102**, 15–17.
- [14] MIURA R 1968 Korteweg-de Vries equation and generalizations. I. A remarkable explicit nonlinear transformation, *J. Math. Phys.* **9** 1202.
- [15] MURYSHEV A, SHLYAPNIKOV G, ERTMER W, SENGSTOCK K, & LEWENSTEIN M 2002 Dynamics of dark solitons in elongated Bose-Einstein condensates, *Phys. Rev. Lett.* **89** 110401.
- [16] RATLIFF D 2017 Phase dynamics of periodic wavetrains leading to the fifth-order KP equation, *Physica D* **353-354** 11–19.
- [17] RATLIFF D 2018 Double degeneracy in multiphase modulation and the emergence of the Boussinesq equation, *Stud. Appl. Math.* **140** 48–77.
- [18] RATLIFF D AND BRIDGES T 2015 Phase dynamics of periodic waves leading to the Kadomtsev-Petviashvili equation in 3+1 dimensions, *Proc. R. Soc. Lond. A* **471** 20150137.
- [19] RATLIFF D AND BRIDGES T 2016 Whitham modulation equations, coalescing characteristics, and dispersive Boussinesq dynamics, *Physica D* **333** 107–116.
- [20] RATLIFF D AND BRIDGES T 2016 Multiphase wavetrains, singular wave interactions, and the emergence of the KdV equation, *Proc. R. Soc. Lond. A* **472** 20160456.
- [21] TUSZYŃSKI J AND DIXON J 1989 Coherent structures in strongly interacting manybody systems. I. Derivation of dynamics, *J. Phys. A Math. Gen.* **22** 4166.
- [22] TUSZYŃSKI J AND KIBLER M 1992 The formation of coherent structures in strongly interacting q-boson systems, *J. Phys. A Math. Gen.* **25** 2425.
- [23] WHITHAM G 1974 *Linear and Nonlinear Waves*, (New York: John Wiley & Sons).
- [24] YUEN H AND LAKE B 1975 Nonlinear deep water waves: Theory and experiment, *Phys. Fluids* **18** 956–960.
- [25] ZHANG D-J, ZHAO S-L, SUN Y-Y, ZHOU J 2014 Solutions to the modified Korteweg-de Vries equation, *Rev. Math. Phys.* **26** 1430006.